arXiv:1505.07236v5 [math.AP] 10 Apr 2016

Self-adjoint elliptic operators with boundary conditions on not closed hypersurfaces

Andrea Mantile[a], Andrea Posilicano[b,*], Mourad Sini[c]

[a]Laboratoire de Mathématiques, Université de Reims - FR3399 CNRS, Moulin de la Housse BP 1039, 51687 Reims, France
[b]DiSAT - Sezione di Matematica, Università dell'Insubria, Via Valleggio 11, I-22100 Como, Italy
[c]RICAM, Austrian Academy of Sciences, Altenbergerstr. 69, A-4040 Linz, Austria

---

Abstract

The theory of self-adjoint extensions of symmetric operators is used to construct self-adjoint realizations of a second-order elliptic differential operator on $\mathbb{R}^n$ with linear boundary conditions on (a relatively open part of) a compact hypersurface. Our approach allows to obtain Kreĭn-like resolvent formulae where the reference operator coincides with the "free" operator with domain $H^2(\mathbb{R}^n)$; this provides an useful tool for the scattering problem from a hypersurface. Concrete examples of this construction are developed in connection with the standard boundary conditions, Dirichlet, Neumann, Robin, δ and δ'-type, assigned either on a $(n-1)$ dimensional compact boundary $\Gamma = \partial\Omega$ or on a relatively open part $\Sigma \subset \Gamma$. Schatten-von Neumann estimates for the difference of the powers of resolvents of the free and the perturbed operators are also proven; these give existence and completeness of the wave operators of the associated scattering systems.

Keywords: Elliptic Operators, Boundary Conditions, Kreĭn's Resolvent Formulae, Self-Adjoint Extensions
2000 MSC: 35J15, 35J25, 47B25, 47F05

---

1. Introduction.

This work is concerned with the self-adjoint realizations of symmetric, second-order elliptic operators

$$Au(x) = \sum_{1\le i,j\le n} \partial_{x_i}(a_{ij}(x)\partial_{x_j}u(x)) - V(x)u(x)\,, \quad x \equiv (x_1,\dots,x_n) \in \mathbb{R}^n\,, \tag{1.1}$$

with boundary conditions on (relatively open parts of) hypersurfaces which are boundaries Γ of bounded open sets $\Omega \subset \mathbb{R}^n$. We assume Ω to be of class $C^{1,1}$; that suffices in case the boundary conditions are globally imposed on Γ whereas, in the case of boundary conditions on $\Sigma \subset \Gamma$, we require more regularity on Γ, even if it suffices to assume Σ to be of class $C^{1,0}$, i.e Σ has a Lipschitz boundary. By [37], [13], we expect that our results can be extended to the case in which also Γ is merely Lipschitz. As regards the conditions on the coefficients a_{ij} and V, for simplicity we assume that they are both in $C^\infty_b(\mathbb{R}^n)$, the standard regularity hypotheses allowing to use the classical results on mapping properties of surface potentials (as given, for example, in [57, Chapter 6]). However our regularity assumptions could possibly be relaxed. For example, in the case $a_{ij} = \delta_{ij}$, it should suffice to work with any $(-\Delta)$-bounded potential V; moreover, by following the results concerning the surface potentials provided in [1] and references therein, we expect that our analysis could also be adapted to the case where the a_{ij}'s are bounded and Lipschitz and V is bounded.

When defined on the domain $\mathrm{dom}(A) = H^2(\mathbb{R}^n)$, the operator A is self-adjoint and bounded from above. We then consider the same differential operator A but now acting on a domain characterized by linear boundary conditions on Γ or on a relatively open part $\Sigma \subset \Gamma$. Using the abstract theory of self-adjoint extensions of symmetric operators developed in [60]-[63], we construct these models as singular perturbations of the "free operator" with domain $H^2(\mathbb{R}^n)$. This allows us to describe all possible linear boundary conditions within an unified framework where the corresponding self-adjoint operators $A_{\Pi,\Theta}$ are parametrized through couples (Π,Θ), where Π is an orthogonal projector on the Hilbert trace space $H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)$ and Θ is a self-adjoint operator in the Hilbert space given by the range of Π. Our approach naturally yields to Kreĭn-type formulae expressing the resolvent of the self-adjoint extension $A_{\Pi,\Theta}$ in terms of the unperturbed resolvent $(-A+z)^{-1}$, plus a

---

[*] Corresponding author

Email addresses: `andrea.mantile@univ-reims.fr` (Andrea Mantile), `andrea.posilicano@uninsubria.it` (Andrea Posilicano), `mourad.sini@oeaw.ac.at` (Mourad Sini)

Preprint submitted to Journal of Differential Equations *April 12, 2016*

non-perturbative term; under suitable regularity assumptions of the parameters (Π,Θ), the difference $(-A_{\Pi,\Theta}+z)^{-k}-(-A+z)^{-k}$ is of trace class (for sufficiently large k) and the Birman-Kato criterion allows to consider $\{A, A_{\Pi,\Theta}\}$ as a scattering system provided with the corresponding wave operators.

Singular perturbations supported on manifolds of lower dimension have been the object of a large number of investigations (see for instance [2]-[5], [8]-[12], [14]-[16], [21]-[33], [35], [36], [42], [51]-[54], [59], [60] and references therein). These have mainly concerned the case of δ-perturbed Schrödinger operators and are generally motivated by the quantum dynamical modelling, as the case of leaky quantum graphs, or the quantum interaction with charged surfaces.

Covering a wider class of models, the analysis developed in our work have been inspired by the scattering problem from a compact hypersurface with abstract boundary conditions. When these conditions are encoded by the extension $A_{\Pi,\Theta}$, the scattered field u_{sc} corresponding to an incident wave u_{in} is expected to be related to a limit absorption principle for $(-A_{\Pi,\Theta}+z)^{-1}$. In particular, the result obtained in the simpler case of point scatterers (see [47]) suggests the relation

$$u_{\mathrm{sc}} = \lim_{\mathbb{C}_+\ni z\to\lambda\in\mathbb{R}} \Big((-A_{\Pi,\Theta}+z)^{-1}(-A+z)u_{\mathrm{in}}\Big) - u_{\mathrm{in}}\,,$$

where the limit is to be understood in an appropriate operator topology. This, using the Kreĭn resolvent identity for $(-A_{\Pi,\Theta}+z)^{-1}-(-A+z)^{-1}$, would lead to an explicit characterization of the scattered field in terms of a factorized formula depending on the incident wave. Different applications of this type of formulas can be foreseen. In the most standard cases (Dirichlet, Neumann and impedance boundary conditions on Γ or $\Sigma\subset\Gamma$), they have been exploited in the analysis of the corresponding inverse scattering problem for surfaces reconstruction (see [50] for an introduction to the factorization method). In this connection, our result could provide an unified method to derive factorized formulas for the scattered field for a large class of scattering problems with rather general linear boundary conditions.

The first part of this work is devoted to the construction of self-adjoint elliptic operators with abstract boundary conditions on Γ. In Section 2 we briefly recall the main results of the extension theory of symmetric operators according to [60]-[63], while, in Section 3 the mapping properties of the trace operators and of the single and double layer operators, related to the surface Γ and to the operator A, are reviewed. In the Section 4, we introduce our model through the symmetric operator S given by the restriction of $A : H^2(\mathbb{R}^n)\subset L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)$, to the dense linear set $\{u\in H^2(\mathbb{R}^n) : u|\Gamma = \partial_{\underline{a}}u|\Gamma = 0\}$, being $\partial_{\underline{a}}$ the co-normal derivative on Γ. The construction of the self-adjoint extensions of S, parametrized through couples (Π,Θ) on the trace space $H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma)$, is then given in Theorem 4.4 and Corollary 4.9, where a Kreĭn-like resolvent formula is also provided. The Schatten-von Neumann type estimates for the difference of the powers of resolvents, together with the spectral properties of these extensions and the existence and completeness of the wave operators are then given in Theorems 4.11, 4.12 and Corollary 4.13.

The second part of the work is devoted to applications. In Section 5 we construct the standard models, i.e. Dirichlet, Neumann, impedance (or Robin), δ and δ'-type boundary conditions on Γ, in terms of extensions of S. The main issue concerns the determination of the parameters (Π,Θ) corresponding to the required constraints. In the case of global conditions on Γ, this task is simplified by the nature of Π, which, in the above mentioned cases, identifies with the projection onto the $H^{\frac{3}{2}}(\Gamma)$ component for the Dirichlet case, with the projection onto the $H^{\frac{1}{2}}(\Gamma)$ component for the Neumann case and with $\Pi = 1$ in the Robin case; then, the determination of Θ for the corresponding boundary conditions easily follows (almost) from algebraic arguments. The case of Dirichlet, Neumann, impedance, δ and δ'-type conditions assigned only on a relatively open subset $\Sigma\subset\Gamma$ is more complex and requires further work: in particular the analysis of self-adjoint operators related to compressions of sesquilinear forms on the subspaces $H^{\frac{3}{2}}(\Sigma)$ and $H^{\frac{1}{2}}(\Sigma)$. This point is developed in Section 6 and in the Appendix. At least to our knowledge, the Kreĭn formulae we provide in the case of boundary conditions on not closed hypersurfaces $\Sigma\subset\Gamma$ do not appear in the past literature.

2. Preliminaries: Self-adjoint extensions of symmetric operators.

Given the self-adjoint operator

$$A : \mathrm{dom}(A)\subseteq\mathscr{H}\to\mathscr{H}$$

in the Hilbert space $\mathscr{H}$ (equipped with the scalar product $\langle\cdot,\cdot\rangle_{\mathscr{H}}$), let

$$\tau : \mathrm{dom}(A)\to\mathfrak{h}$$

be continuous (w.r.t. the graph norm in $\mathrm{dom}(A)$) and surjective onto the auxiliary Hilbert space $\mathfrak{h}$ (equipped with the scalar product $\langle\cdot,\cdot\rangle_{\mathfrak{h}}$). We further assume that $\ker(\tau)$ is dense in $\mathscr{H}$ and introduce the densely defined, closed, symmetric operator

$$S := A|\ker(\tau)\,.$$

Our aim is to provide all self-adjoint extensions of S; here we use the approach developed in [60]-[63] to which we refer for proofs and for the connections with other well known approaches to this problem (von Neumann's theory and boundary triples theory) .

For notational convenience we do not identify $\mathfrak{h}$ with its dual $\mathfrak{h}'$ and we denote by $J : \mathfrak{h} \to \mathfrak{h}'$ the duality mapping (a bijective isometry) given by the canonical isomorphism from $\mathfrak{h}$ onto $\mathfrak{h}'$, i.e. $J(\varphi)$ is the differential of the function $\varphi \mapsto \frac{1}{2}\,\|\varphi\|^2_{\mathfrak{h}}$ (see e.g. [6, Section 3.1]); $\mathfrak{h}'$ inherits a Hilbert space structure by the scalar product $\langle \phi_1, \phi_2\rangle_{\mathfrak{h}'} := \langle J^{-1}\phi_1, J^{-1}\phi_2\rangle_{\mathfrak{h}}$, so that J becomes an unitary map, and we denote by $\langle\cdot,\cdot\rangle_{\mathfrak{h}'\mathfrak{h}}$ the $\mathfrak{h}'$-$\mathfrak{h}$ duality

$$\langle \phi, \varphi\rangle_{\mathfrak{h}'\mathfrak{h}} := \langle J^{-1}\phi, \varphi\rangle_{\mathfrak{h}}\,.$$

The inverse $J^{-1} : \mathfrak{h}' \to \mathfrak{h}$ gives the duality mapping from $\mathfrak{h}'$ to its dual $\mathfrak{h}'' \equiv \mathfrak{h}$; we denote by $\langle\cdot,\cdot\rangle_{\mathfrak{h}\mathfrak{h}'}$ the $\mathfrak{h}$-$\mathfrak{h}'$ duality defined by $\langle \varphi, \phi\rangle_{\mathfrak{h}\mathfrak{h}'} := \langle J\varphi, \phi\rangle_{\mathfrak{h}'} = \langle \varphi, J^{-1}\phi\rangle_{\mathfrak{h}}$.

Given a densely defined linear operator

$$\Xi : \mathrm{dom}(\Xi) \subseteq \mathfrak{h} \to \mathfrak{h}$$

we denote by Ξ' the dual operator

$$\Xi' : \mathrm{dom}(\Xi') \subseteq \mathfrak{h}' \to \mathfrak{h}'\,, \qquad \Xi\phi := \phi'\,,$$

$$\mathrm{dom}(\Xi') := \{\phi \in \mathfrak{h}' : \exists \phi' \in \mathfrak{h}' \text{ such that } \langle \phi', \varphi\rangle_{\mathfrak{h}'\mathfrak{h}} = \langle \phi, \Xi\varphi\rangle_{\mathfrak{h}'\mathfrak{h}} \text{ for all } \varphi \in \mathrm{dom}(\Xi)\}\,;$$

this is related to the (Hilbert) adjoint $\Xi^* : \mathrm{dom}(\Xi^*) \subseteq \mathfrak{h} \to \mathfrak{h}$ by

$$\Xi^* = J^{-1}\Xi' J\,, \qquad \mathrm{dom}(\Xi^*) = J^{-1}(\mathrm{dom}(\Xi'))\,.$$

In the case $\Xi : \mathrm{dom}(\Xi) \subseteq \mathfrak{h}' \to \mathfrak{h}$, the dual operator is defined in a similar way:

$$\Xi' : \mathrm{dom}(\Xi') \subseteq \mathfrak{h}' \to \mathfrak{h}\,, \qquad \Xi'\phi := \varphi\,,$$

$$\mathrm{dom}(\Xi') := \{\phi \in \mathfrak{h}' : \exists \varphi \in \mathfrak{h} \text{ such that } \langle \varphi, \psi\rangle_{\mathfrak{h}\mathfrak{h}'} = \langle \phi, \Xi\psi\rangle_{\mathfrak{h}'\mathfrak{h}} \text{ for all } \psi \in \mathrm{dom}(\Xi)\}\,,$$

or, equivalently,

$$\Xi' : \mathrm{dom}(\Xi') \subseteq \mathfrak{h}' \to \mathfrak{h}\,, \qquad \Xi'\phi := \varphi\,,$$

$$\mathrm{dom}(\Xi') := \{\phi \in \mathfrak{h}' : \exists \varphi \in \mathfrak{h} \text{ such that } \langle \varphi, J^{-1}\psi\rangle_{\mathfrak{h}} = \langle J^{-1}\phi, \Xi\psi\rangle_{\mathfrak{h}} \text{ for all } \psi \in \mathrm{dom}(\Xi)\}\,.$$

The latter definition shows that $\Xi = \Xi'$ if and only if $\tilde{\Xi} := \Xi J : J^{-1}(\mathrm{dom}(\Xi)) \subseteq \mathfrak{h} \to \mathfrak{h}$ is self-adjoint, i.e $\tilde{\Xi}^* = \tilde{\Xi}$; by a slight abuse of terminology we say that Ξ is self-adjoint (resp. symmetric) whenever $\Xi = \Xi'$ (resp. $\Xi \subset \Xi'$).

For any $z \in \rho(A)$ we define $R_z \in \mathsf{B}(\mathscr{H}, \mathrm{dom}(A))$ and $G_z \in \mathsf{B}(\mathfrak{h}', \mathscr{H})$ by

$$R_z := (-A + z)^{-1}\,, \qquad G_z : \mathfrak{h}' \to \mathscr{H}\,, \quad G_z := (\tau R_{\bar{z}})'\,,$$

i.e.

$$\forall \phi \in \mathfrak{h}'\,, \forall u \in \mathscr{H}\,, \quad \langle G_z\phi, u\rangle_{\mathscr{H}} = \langle \phi, \tau(-A + \bar{z})^{-1}u\rangle_{\mathfrak{h}'\mathfrak{h}}\,. \tag{2.1}$$

By our hypotheses on the map τ one gets (see [60, Remark 2.9])

$$\mathrm{ran}(G_z) \cap \mathrm{dom}(A) = \{0\}\,, \tag{2.2}$$

and (see [60, Lemma 2.1])

$$(z - w)\, R_w G_z = G_w - G_z\,, \tag{2.3}$$

so that

$$\mathrm{ran}(G_w - G_z) \subseteq \mathrm{dom}(A) \tag{2.4}$$

and

$$A(G_z - G_w) = z\, G_z - w\, G_w\,. \tag{2.5}$$

Now, in order to simplify the exposition and since such an hypothesis holds true in the applications further considered, we suppose that A has a spectral gap, i.e.

$$\rho(A) \cap \mathbb{R} \neq \emptyset\,.$$

Then we pose

$$G := G_{\lambda_\circ}\,,\quad \lambda_\circ \in \rho(A)\cap\mathbb{R}$$

and define

$$M_z := \tau(G - G_z) \equiv (z-\lambda_\circ)G'G_z : \mathfrak{h}' \to \mathfrak{h}\,. \tag{2.6}$$

Given an orthogonal projection $\Pi : \mathfrak{h} \to \mathfrak{h}$, the dual map $\Pi' : \mathfrak{h}' \to \mathfrak{h}'$ is an orthogonal projection in $\mathfrak{h}'$ (since $\Pi' = J\Pi J^{-1}$ and J is unitary) and, by [6, Proposition 3.5.1], $\operatorname{ran}(\Pi)' = \mathfrak{h}'/\operatorname{ran}(\Pi')^\perp = \operatorname{ran}(\Pi')$ and $\operatorname{ran}(\Pi')' = \mathfrak{h}/\operatorname{ran}(\Pi)^\perp = \operatorname{ran}(\Pi)$. Thus, for a densely defined linear map $\Xi : \operatorname{dom}(\Xi) \subseteq \operatorname{ran}(\Pi') \to \operatorname{ran}(\Pi)$, one has $\Xi' : \operatorname{dom}(\Xi') \subseteq \operatorname{ran}(\Pi') \to \operatorname{ran}(\Pi)$. As in the case $\operatorname{ran}(\Pi) = \mathfrak{h}$ we say that Ξ is symmetric whenever $\Xi \subset \Xi'$ and that is self-adjoint whenever $\Xi = \Xi^{'}$; one has that $\Xi = \Xi'$ (resp. $\Xi \subseteq \Xi'$) if and only if $\tilde\Xi = \tilde\Xi^*$ (resp. $\tilde\Xi \subseteq \tilde\Xi^*$), where $\tilde\Xi := \Xi J$, $\operatorname{dom}(\tilde\Xi) := J^{-1}(\operatorname{dom}(\Xi))$. Finally, given a self-adjoint $\Theta : \operatorname{dom}(\Theta) \subseteq \operatorname{ran}(\Pi') \to \operatorname{ran}(\Pi)$, we define

$$Z_{\Pi,\Theta} := \{z \in \rho(A) : \Theta + \Pi M_z \Pi' \text{ has a bounded inverse on } \operatorname{ran}(\Pi) \text{ to } \operatorname{ran}(\Pi')\}\,.$$

Theorem 2.1. *Any self-adjoint extension of* $S = A|\ker(\tau)$ *is of the kind* $A_{\Pi,\Theta}$, *where* $\Pi : \mathfrak{h} \to \mathfrak{h}$ *is an orthogonal projection,* $\Theta : \operatorname{dom}(\Theta) \subseteq \operatorname{ran}(\Pi') \to \operatorname{ran}(\Pi)$ *is a self-adjoint operator and*

$$\operatorname{dom}(A_{\Pi,\Theta}) := \{u = u_\circ + G\phi\,,\ u_\circ \in \operatorname{dom}(A)\,,\ \phi \in \operatorname{dom}(\Theta)\,,\ \Pi\tau u_\circ = \Theta\phi\}\,,$$

$$A_{\Pi,\Theta}u := Au_\circ + \lambda_\circ G\phi\,.$$

Moreover $Z_{\Pi,\Theta}$ *is not void,* $\mathbb{C}\backslash\mathbb{R} \subseteq Z_{\Pi,\Theta} \subseteq \rho(A_{\Pi,\Theta})$, *and the resolvent of the self-adjoint extension* $A_{\Pi,\Theta}$ *is given by the Kreĭn's type formula*

$$(-A_{\Pi,\Theta} + z)^{-1} = R_z + G_z\Pi'(\Theta + \Pi M_z\Pi')^{-1}\Pi\tau R_z\,,\quad z \in Z_{\Pi,\Theta}\,. \tag{2.7}$$

Proof. Let us pose $\tilde\Theta := \Theta J : J^{-1}(\operatorname{dom}(\Theta)) \subseteq \operatorname{ran}(\Pi) \to \operatorname{ran}(\Pi)$, $\tilde G_z := G_z J : \mathfrak{h} \to \mathscr{H}$ and $\tilde M_z := M_z J : \mathfrak{h} \to \mathfrak{h}$. Thus $\tilde\Theta$ is self-adjoint in $\operatorname{ran}(\Pi)$,

$$\begin{aligned} G_z\Pi'(\Theta + \Pi M_z\Pi')^{-1}\Pi\tau R_z =& G_zJ\Pi J^{-1}((\Theta J + \Pi\tilde M_z\Pi)J^{-1})^{-1}\Pi\tau R_z \\ =& \tilde G_z\Pi(\tilde\Theta + \Pi\tilde M_z\Pi)^{-1}\Pi\tau G_z \end{aligned}$$

and $Z_{\Pi,\Theta} = \{z \in \rho(A) : 0 \in \rho(\tilde\Theta + \Pi\tilde M_z\Pi)\}$. Therefore, by [63, Theorem 2.1] (see Remark 2.2 below), the linear operator

$$\hat A_{\Pi,\Theta} : \operatorname{dom}(\tilde A_{\Pi,\Theta}) \subseteq \mathscr{H} \to \mathscr{H}\,,\quad (-\hat A_{\Pi,\Theta} + z) := (-A + z)u_z\,,$$

$$\operatorname{dom}(\hat A_{\Pi,\Theta}) = \{u = u_z + G_z\Pi'(\Theta + \Pi M_z\Pi')^{-1}\Pi\tau u_z,\ u_z \in \operatorname{dom}(A)\,,\ z \in Z_{\Pi,\Theta}\}$$

is a z-independent self-adjoint extension of $A|\ker(\tau)$; moreover its resolvent is given by (2.7). Let us now show that $\hat A_{\Pi,\Theta} = A_{\Pi,\Theta}$. At first we pose $\phi_z := (\Theta + \Pi M_z\Pi')^{-1}\Pi\tau u_z$, so that, since the definition of $\hat A_{\Pi,\Theta}$ is z-independent, $u \in \operatorname{dom}(\hat A_\Theta)$ if and only if for any $z \in Z_{\Pi,\Theta}$ there exists $u_z \in \operatorname{dom}(A_\circ)$, $\Pi\tau u_z = (\Theta + \Pi M_z\Pi)\phi_z$, such that $u = u_z + G_z\phi_z$. Therefore, by (2.3),

$$u_z - u_w = G_w\phi_w - G_z\phi_z = G_z(\phi_w - \phi_z) + (z - w)R_zG_w\phi_w\,.$$

By (2.2), one obtains $G_z(\phi_w - \phi_z) = 0$. Since G_z is injective (it is the adjoint of a surjective map), this gives $\phi_z = \phi_w$, i.e. the definition of ϕ_z is z-independent. Thus, posing $u_\circ := u_z + (G_z - G)\phi$, one has $u = u_\circ + G\phi$, with $u_\circ \in \operatorname{dom}(A)$ and

$$\Pi\tau u_\circ = \Pi\tau u_z + \Pi\tau(G_z - G)\phi = (\Theta + \Pi M_z\Pi)\phi - \Pi M_z\Pi\phi = \Theta\phi\,.$$

Then, by (2.5),

$$A_{\Pi,\Theta}u = Au_z + zG_z\phi = Au_\circ - A(G_z - G)\phi + zG_z\phi = Au_\circ + \lambda_\circ G\phi\,.$$

Finally, by [63, Corollary 3.2] (also see [61, Theorem 4.3]), any self-adjoint extension of $A|\ker(\tau)$ is of the kind $A_{\Pi,\Theta}$ for some couple (Π, Θ). ☐

Remark 2.2. Let us notice that the operators denoted by G_z and Γ_z in [62] and [63] here correspond to $\tilde G_z$ and $\tilde M_z$ respectively.

Let us remark that we have not used neither the adjoint S^* nor the defect space $\ker(S^* - z)$. However these can be readily obtained:

Lemma 2.3.

$$\mathrm{dom}(S^*) = \{u = u_\circ + G\phi\,,\ u_\circ \in \mathrm{dom}(A)\,,\ \phi \in \mathfrak{h}'\}\,,$$
$$S^*u = Au_\circ + \lambda_\circ G\phi\,,$$

and

$$\ker(S^* - z) = \{G_z\phi\,,\ \phi \in \mathfrak{h}'\}\,, \quad z \in \rho(A)\,.$$

Proof. By [62, Theorem 3.1] (see Remark 2.2) one has

$$\mathrm{dom}(S^*) = \{u = u_* + \frac{1}{2}\,(G_i + G_{-i})\phi\,,\ u_* \in \mathrm{dom}(A)\,,\ \phi \in \mathfrak{h}'\}$$

$$S^*u = Au_* + \frac{i}{2}\,(G_i - G_{-i})\phi\,.$$

Thus, posing $u_\circ := u_* + \frac{1}{2}\,(G_i - G)\phi + \frac{1}{2}\,(G_{-i} - G)\phi$, one has

$$\mathrm{dom}(S^*) = \{u = u_\circ + G\phi\,,\ u_\circ \in \mathrm{dom}(A)\,,\ \phi \in \mathfrak{h}'\}$$

and

$$S^*u = Au_\circ - \frac{1}{2}\,A(G_i - G)\phi - \frac{1}{2}\,A(G_{-i} - G)\phi + \frac{i}{2}\,(G_i - G_{-i})\phi\,.$$

By (2.5) one then obtains $S^*u = Au_\circ + \lambda_\circ G\phi$.

The vector $u = u_\circ + G\phi \in \mathrm{dom}(S^*)$ belong to $\ker(S^* - z)$ if and only if $(\lambda_\circ - z)G\phi = (-A + z)u_\circ$. This gives $u = (\lambda_\circ - z)R_zG\phi + G\phi$; by (2.3) one gets $u = G_z\phi$. ☐

Remark 2.4. By Lemma 2.3 and Theorem 2.1, any self-adjoint extension of S is of the kind

$$A_{\Pi,\Theta} := S^*|\mathrm{dom}(A_{\Pi,\Theta})\,,$$

$$\mathrm{dom}(A_{\Pi,\Theta}) = \{u \in \mathrm{dom}(S^*) : \beta_0 u \in \mathrm{dom}(\Theta)\,,\ \Pi\beta_1 u = \Theta\beta_0 u\}\,,$$

where

$$\beta_0 : \mathrm{dom}(S^*) \to \mathfrak{h}'\,, \quad \beta_0 u := \phi\,.$$
$$\beta_1 : \mathrm{dom}(S^*) \to \mathfrak{h}\,, \quad \beta_1 u := \tau u_\circ\,,$$

Moreover (using [62, Theorem 3.1]) one has the abstract Green's identity

$$\langle S^*u, v\rangle_{\mathscr{H}} - \langle u, S^*v\rangle_{\mathscr{H}} = \langle \beta_1 u, \beta_0 v\rangle_{\mathfrak{h},\mathfrak{h}'} - \langle \beta_0 u, \beta_1 v\rangle_{\mathfrak{h}',\mathfrak{h}}$$

Let us also notice that $u = G_z\phi$ solves the adjoint (abstract) boundary value problem

$$\begin{cases} S^*u = z\,u \\ \beta_0 u = \phi\,. \end{cases} \tag{2.8}$$

Remark 2.5. By [62, Theorem 3.1], the triple $(\mathfrak{h}, \Gamma_1, \Gamma_2)$, where $\Gamma_1 := -J^{-1}\beta_0$ and $\Gamma_2 := \beta_1$ is a boundary triplet for S^*, i.e. Γ_1 and Γ_2 are surjective and

$$\langle S^*u, v\rangle_{\mathscr{H}} - \langle u, S^*v\rangle_{\mathscr{H}} = \langle \Gamma_1 u, \Gamma_2 v\rangle_{\mathfrak{h}} - \langle \Gamma_2 u, \Gamma_1 v\rangle_{\mathfrak{h}}$$

holds true. The Weyl function of the boundary triple $(\mathfrak{h}, \Gamma_1, \Gamma_1)$ is the bounded linear operator $M_zJ : \mathfrak{h} \to \mathfrak{h}$, where M_z is defined in (2.6) (see [62, Theorem 3.1]). For Boundary Triple Theory we refer to [20], [65] and references therein.

We conclude the section with the following result:

Lemma 2.6. *Given the linear operator* $\Xi : \mathrm{dom}(\Xi) \subseteq \mathrm{ran}(\Pi') \to \mathrm{ran}(\Pi)$, *let us define the linear operator* $A_{\Pi,\Xi}$ *by*

$$A_{\Pi,\Xi} := S^*|\mathrm{dom}(A_{\Pi,\Xi})\,, \quad \mathrm{dom}(A_{\Pi,\Xi}) := \{u \in \mathrm{dom}(S^*) : \beta_0 u \in \mathrm{dom}(\Xi)\,,\ \Pi\beta_1 u = \Xi\beta_0 u\}\,.$$

Then $A_{\Pi,\Xi}$ *is self-adjoint if and only if* Ξ *is self-adjoint.*

Proof. Since we can re-write $\mathrm{dom}(A_{\Pi,\Xi})$ as

$$\mathrm{dom}(A_{\Pi,\Xi}) = \{u \in \mathrm{dom}(S^*) : \Gamma_1 u \in J^{-1}(\mathrm{dom}(\Xi))\,,\ \Pi\Gamma_2 u = -\Xi J\Gamma_1 u\}\,,$$

and since $(\mathfrak{h}, \Gamma_1, \Gamma_2)$ is a boundary triple, by e.g. [65, Lemma 14.6 and Theorem 14.7], $A_{\Pi,\Xi}$ is self-adjoint if and only if $\Xi J : J^{-1}\mathrm{dom}(\Xi) \subseteq \mathrm{ran}(\Pi) \to \mathrm{ran}(\Pi)$ is self-adjoint. The latter is equivalent to Ξ self-adjoint. ☐

3. Preliminaries: Sobolev spaces and boundary-layer operators

3.1. Sobolev spaces.

Let Ω be a non-empty open subset of $\mathbb{R}^n$; $H^k(\Omega)$, $k \in \mathbb{N}$, denotes the usual Sobolev-Hilbert spaces $H^k(\Omega) := \{u \in L^2(\Omega) : \partial^\alpha u \in L^2(\Omega)\,,\ |\alpha| \le k\}$, where $\partial^\alpha : \mathscr{D}'(\Omega) \to \mathscr{D}'(\Omega)$ denotes the distributional partial derivatives of order $|\alpha|$. In the case $\Omega = \mathbb{R}^n$, the scale of Sobolev-Hilbert spaces $H^s(\mathbb{R}^n)$, $s \in \mathbb{R}$, is defined by $H^s(\mathbb{R}^n) := \{u \in \mathscr{S}'(\mathbb{R}^n) : \int_{\mathbb{R}^n} |\hat{u}(\kappa)|^2(|\kappa|^2+1)^s d\kappa < +\infty\}$, where $\mathscr{S}'(\mathbb{R}^n)$ is the space of tempered distributions and $\hat{u}$ denotes Fourier's transform. $H^{-s}(\mathbb{R}^n)$ identifies with the dual space of $H^s(\mathbb{R}^n)$; we denote by $\langle\cdot,\cdot\rangle_{-s,s}$ the H^{-s}-H^s duality pairing.

Let us now suppose that Ω is bounded and let Γ denote its boundary: $\Gamma = \partial\Omega$. We further suppose that $\Omega \subset \mathbb{R}^n$ is of class $C^{k,1}$, $k \ge 0$, i.e we suppose that its boundary Γ is a manifold of dimension $n-1$ whose local maps are Lipschitz-continuous, together with their inverses, up to the order k. In the particular case $C^{0,1}$, Ω is referred to as a Lipschitz domain. The scale of Sobolev-Hilbert space $H^s(\Omega)$, $s \in \mathbb{R}$, is then defined by $H^s(\Omega) := \{u|\Omega : u \in H^s(\mathbb{R}^n)\}$, $u|\Omega$ denoting the restriction of u to Ω. In the case $s \in \mathbb{N}$, this definition reproduces the previous one (see e.g. [39, 1.4.3.1]).

The Sobolev spaces of L^2-functions on Γ, next denoted with $H^s(\Gamma)$, are defined by using an atlas of Γ and the Sobolev space on flat, open, bounded, $(n-1)$-dimensional domains (see e.g. [39, Section 1.3.3], [57, Chapter 3], [67, Section 3.1]); they are well defined up to the order $|s| = k+1$, and $H^{-s}(\Gamma)$ identifies with the dual space of $H^s(\Gamma)$. We denote by $\langle\cdot,\cdot\rangle_{-s,s}$ the H^{-s}-H^s duality pairing. If Γ is a C^2 manifold, considering the Riemannian structure inherited from $\mathbb{R}^n$, we have that $H^s(\Gamma)$ identifies with $\mathrm{dom}((-\Delta_\Gamma)^{\frac{s}{2}})$ with respect of the scalar product

$$\langle\phi,\varphi\rangle_{H^s(\Gamma)} := \langle\Lambda^s\phi,\Lambda^s\varphi\rangle_{L^2(\Gamma)}\,,\qquad \Lambda := (-\Delta_\Gamma+1)^{\frac{1}{2}}\,, \tag{3.1}$$

being Δ_Γ the self-adjoint operator in $L^2(\Gamma)$ corresponding to the Laplace-Beltrami operator on the complete Riemannian manifold Γ (see e.g. [58, Remark 7.6, Chapter 1]). According to this definition, Λ^r is self-adjoint in $H^s(\Gamma)$ with domain $H^{s+r}(\Gamma)$ and acts as a unitary map $\Lambda^r : H^s(\Gamma) \to H^{s-r}(\Gamma)$. In particular $\Lambda^{2s} : H^s(\Gamma) \to H^{-s}(\Gamma)$ plays the role of the duality mapping J introduced in Section 2, and one has $\langle\phi,\psi\rangle_{-s,s} = \langle\Lambda^{-s}\phi,\Lambda^s\psi\rangle_{L^2(\Gamma)}$. In the case Γ is not C^2 one can use the definition (which works in the case Ω is of class $C^{0,1}$) $\Delta_\Gamma := \mathrm{Div}\circ\nabla_{\tan}$ provided in [38, Theorem 1.2]. The two definitions coincide in the case Γ is C^2 (see [38, Section 7]).

Denoting by $0 = \lambda_0 < \lambda_1 \le \cdots \le \lambda_k \le \lambda_{k+1} \le \ldots$, the increasing sequence of the eigenvalues of the self-adjoint operator $-\Delta_\Gamma : \mathrm{dom}(-\Delta_\Gamma) \subset L^2(\Gamma) \to L^2(\Gamma)$ and by $\{\varphi_k\}_{k=1}^\infty$ the corresponding normalized eigenfunctions, one has

$$\|\phi\|^2_{H^s(\Gamma)} = \sum_{k=0}^\infty (\lambda_k+1)^s|\langle\phi,\varphi_k\rangle_{L^2(\Gamma)}|^2\,.$$

Thus, given $r < s < t$, for any $\epsilon > 0$ there exists $c_\epsilon > 0$, $c_\epsilon \uparrow \infty$ as $\epsilon \downarrow 0$, such that

$$\|\phi\|_{H^s(\Gamma)} \le \epsilon\,\|\phi\|_{H^t(\Gamma)} + c_\epsilon\|\phi\|_{H^r(\Gamma)}\,. \tag{3.2}$$

If $\Sigma \subset \Gamma$ is relatively open, then $H^s(\Sigma)$, $|s| \le k+1$, is constructed using an atlas of Σ and the Sobolev spaces on flat $(n-1)$-dimensional domains. If Σ is of class $C^{0,1}$, i.e. if its boundary is a Lipschitz manifold, then a continuation map allows the identification $H^s(\Sigma) = \{\phi|\Sigma : \phi \in H^s(\Gamma)\}$ and $\|\varphi\|_{H^s(\Sigma)} = \inf\{\,\|\phi\|_{H^s(\Gamma)} : \varphi = \phi|\Sigma\}$ (see e.g. [39, Theorem 1.4.3.1], [17, Definition 3.6], [46, Section 4.3]).

In the sequel, we shall also use some closed subspaces of $H^s(\Gamma)$: let $X \subset \Gamma$ denote either $\Sigma^c \equiv \Gamma\backslash\Sigma$ or $\overline{\Sigma}$ and $s \ge 0$, we define

$$H^s_X(\Gamma) := \{\phi \in H^s(\Gamma) : \mathrm{supp}(\phi) \subseteq X\} \equiv H^s(\Gamma) \cap L^2_X(\Gamma)\,,$$

where $L^2_X(\Gamma)$ denotes the space of square-integrable functions with essential support contained in X, and

$$H^{-s}_X(\Gamma) := \{\phi \in H^{-s}(\Gamma) : \langle\phi,\psi\rangle_{-s,s} = 0,\ \text{for any } \psi \in H^s_{\overline{X^c}}(\Gamma)\}\,.$$

Therefore

$$\forall\phi \in H^{-s}_X(\Gamma),\ \forall\varphi \in H^s_{\overline{X^c}}(\Gamma)\,,\qquad \langle\phi,\varphi\rangle_{-s,s} = 0\,. \tag{3.3}$$

By the continuos (with dense range) embeddings $H^s(\Gamma) \hookrightarrow H^r(\Gamma)$, $r < s$, one gets

$$H^r_X(\Gamma) \cap H^s(\Gamma) = H^s_X(\Gamma)\,,\quad H^s_X(\Gamma) \subset H^r_X(\Gamma)\ \text{(dense inclusion)}\,.$$

One has the identifications

$$H^s(\Sigma)' \simeq H^{-s}_{\overline{\Sigma}}(\Gamma)\,, \qquad H^s_{\overline{\Sigma}}(\Gamma)' \simeq H^{-s}(\Sigma)$$

(see [17, Proposition 3.5 and remarks at page 111], [46, Lemma 4.3.1]) and the (strict, whenever $s \ge \frac{1}{2}$) inclusions

$$H^s_{\overline{\Sigma}}(\Gamma) \subseteq H^s(\Sigma) \subseteq L^2(\Sigma) \subseteq H^{-s}(\Sigma) \subseteq H^{-s}_{\overline{\Sigma}}(\Gamma)\,, \quad s > 0\,.$$

Denoting by $H^s_0(\Sigma)$, $s > 0$, the completion of $H^s_{\overline{\Sigma}}(\Gamma)$ with respect to the norm of $H^s(\Sigma)$, one has (see [46, relation (4.3.10)])

$$H^s_{\overline{\Sigma}}(\Gamma) = \begin{cases} H^s_0(\Sigma) & s \neq [s] + \frac{1}{2} \\ H^s_{00}(\Sigma) & s = [s] + \frac{1}{2}\,, \end{cases}$$

where

$$H^s_{00}(\Sigma) := \{\phi \in H^s_0(\Sigma) : d^{-\frac{1}{2}} D^\alpha \phi \in L^2(\Sigma)\,, |\alpha| = [s]\}\,,$$

d denotes the distance to the boundary $\partial\Sigma$ and D denotes the covariant derivative. In particular, since $H^{\frac{1}{2}}_0(\Sigma) = H^{\frac{1}{2}}(\Sigma)$, it results

$$\phi \in H^{\frac{1}{2}}(\Sigma) \quad \text{and} \quad \int_\Sigma \frac{|\phi(x)|^2}{d(x)}\, d\sigma_\Gamma(x) < +\infty \qquad \Longleftrightarrow \qquad \tilde\phi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)\,,$$

where σ_Γ denotes the surface measure and here $\tilde\phi$ denotes the extension by zero.

We shall also need the Hilbert orthogonal

$$H^s_{\Sigma^c}(\Gamma)^\perp = \{\phi \in H^s(\Gamma) : \langle \Lambda^{2s}\phi, \psi\rangle_{-s,s} = 0,\ \text{for any } \psi \in H^s_{\Sigma^c}(\Gamma)\} = \Lambda^{-2s} H^{-s}_{\overline{\Sigma}}(\Gamma)\,. \tag{3.4}$$

Let $\Pi_\Sigma : H^s(\Gamma) \to H^s(\Gamma)$ be the orthogonal projection onto $H^s_{\Sigma^c}(\Gamma)^\perp$, then (see e.g. [57, page 77]) the map

$$U_\Sigma : H^s_{\Sigma^c}(\Gamma)^\perp \to H^s(\Sigma)\,, \quad \mathrm{U}_\Sigma(\Pi_\Sigma\phi) := (\Pi_\Sigma\phi)|\Sigma = \phi|\Sigma$$

is an unitary isomorphism. Therefore we can regard $H^s(\Sigma)$ as a closed subspace of $H^s(\Gamma)$. Using the decomposition $\phi = (1 - \Pi_\Sigma)\phi \oplus U_\Sigma^{-1}(\phi|\Sigma)$, the restriction operator $R_\Sigma\phi := 0 \oplus \mathrm{U}_\Sigma\Pi_\Sigma\phi = 0 \oplus (\phi|\Sigma)$ is the orthogonal projection from $H^s(\Gamma) \simeq H^s_{\Sigma^c}(\Gamma) \oplus H^s(\Sigma)$ onto $H^s(\Sigma)$.

3.2. Sobolev Multipliers

Let us now introduce the following notation: we write $\psi \in M^s(\Gamma)$, $s \ge 0$, whenever ψ is a multiplier in $H^s(\Gamma)$, i.e.

$$\begin{aligned} M^s(\Gamma) :=& \{\psi \in H^s(\Gamma) : \psi\phi \in H^s(\Gamma) \text{ for any } \phi \in H^s(\Gamma)\} \\ =&\{\psi \in H^s(\Gamma) : \exists\, m_\psi \ge 0 \text{ s.t. } \|\psi\phi\|_{H^s(\Gamma)} \le m_\psi\, \|\phi\|_{H^s(\Gamma)}\}\,. \end{aligned} \tag{3.5}$$

The equality holds, by the closed graph theorem, since the map $\phi \mapsto \psi\phi$ is closed and everywhere defined. Notice that $M^s(\Gamma) \subseteq L^\infty(\Gamma)$:

$$\|\psi\|_{L^\infty(\Gamma)} = \lim_{k\to+\infty} \|\psi\|_{L^{2k}(\Gamma)} = \lim_{k\to+\infty} \|\psi^k\|^{1/k}_{L^2(\Gamma)} \le \lim_{k\to+\infty} \|\psi^k\|^{1/k}_{H^s(\Gamma)} \le m_\psi \lim_{k\to+\infty} \|1\|^{1/k}_{H^s(\Gamma)} = m_\psi\,.$$

By the same kind of proofs which hold in the flat case (see [56, Proposition 3.5.1, Corollary 3.5.7]) one has

$$M^s(\Gamma) \subseteq M^r(\Gamma)\,, \quad r \le s\,,$$

and

$$\psi \in M^s(\Gamma) \text{ and } 1/\psi \in L^\infty(\Gamma) \quad \Rightarrow \quad 1/\psi \in M^s(\Gamma)\,.$$

We recall some relatively simple sufficient conditions in order that a given function belongs to $M^s(\Gamma)$. By [57, Theorem 3.20],

$$W^{k,\infty}(\Gamma) \subseteq M^s(\Gamma)\,, \qquad k \ge \max\{1, s\}\,,$$

where $W^{k,\infty}(\Gamma)$ denotes the set of functions in $C^{k-1}(\Gamma)$ with k-order distributional derivatives in $L^\infty(\Gamma)$. By [67, Proposition 4.5] this result can be improved to

$$A^s(\Gamma) \subseteq M^s(\Gamma)\,,$$

where (see [67, Section 4.1]) $A^k(\Gamma) = W^{k,\infty}(\Gamma)$ whenever $s = k$ and, in case $s > 0$ is not an integer,

$$A^s(\Gamma) := \left\{\phi \in W^{[s],\infty}(\Gamma) : q_{i,\lambda,\phi} \in L^\infty(f_i(U_i))\,,\ i \in I\,,\ \lambda = s - [s]\right\}\,,$$

$$q_{\phi,\lambda,i}(x) := \sum_{|\alpha|\le[s]} \int_{f_i(U_i)} \frac{|\partial^\alpha\phi_i(x) - \partial^\alpha\phi_i(y)|^2}{\|x-y\|^{n-1+2\lambda}}\,dy\,, \quad \phi_i := (\varphi_i\phi)\circ f_i^{-1}\,.$$

Here $\{(U_i, f_i)\}_{i\in I}$ is an admissible atlas of Γ and $\{\varphi_i\}_{i\in I}$ is a subordinate partition of unity.

Notice that, for any $s-[s] < \kappa < 1$, one has $\tilde{C}^{[s],\kappa}(\Gamma) \subseteq A^s(\Gamma)$, where $\tilde{C}^{k,\kappa}(\Gamma)$ denotes the set of functions in $W^{k,\infty}(\Gamma)$ having Hölder continuos (of exponent κ) k-order derivatives.

In the case Γ is a smooth manifold, by [19, Theorem 24] one gets

$$H^s(\Gamma) = M^s(\Gamma)\,, \qquad s > \frac{1}{2}(n-1)\,.$$

By the embedding $H^s(\Gamma) \hookrightarrow L^q(\Gamma)$, $\frac{1}{q} = \frac{1}{2} - \frac{s}{n-1}$, which holds whenever $s < \frac{1}{2}(n-1)$, and by [19, Theorem 27], one gets

$$L^{(n-1)/s}_s(\Gamma) \cap L^\infty(\Gamma) \subseteq M^s(\Gamma)\,, \qquad s < \frac{1}{2}(n-1)\,,$$

where

$$L^q_s(\Gamma) := \{\phi \in L^q(\Gamma) : (-\Delta_\Gamma)^{\frac{s}{2}}\phi \in L^q(\Gamma)\}\,.$$

3.3. *Trace maps.*

For a bounded open domain Ω of class $C^{k,1}$, we pose

$$\Omega_+ := \mathbb{R}^n\backslash\overline{\Omega}\,, \qquad \Omega_- := \Omega\,,$$

while ν denotes the outward normal vector on Γ. The one-sided, zero-order, trace operators $\gamma_0^\pm$ act on a smooth function $u \in C^\infty\left(\overline{\Omega_\pm}\right)$ as $\gamma_0^\pm u = u|\Gamma$, where $\varphi|\Gamma$ is the restriction to Γ. These maps uniquely extend to bounded linear operators (see e.g. [57, Theorem 3.37])

$$\gamma_0^\pm \in \mathsf{B}(H^s(\Omega_\pm), H^{s-\frac{1}{2}}(\Gamma))\,, \qquad \frac{1}{2} < s \le k+1\,. \tag{3.6}$$

Then, given $a_{ij} \in C^\infty(\mathbb{R}^n)$, $a_{ij}(x) = a_{ji}(x)$ such that

$$\forall x,\xi \in \mathbb{R}^n\,, \quad \sum_{1\le i,j\le n} a_{ij}(x)\xi_i\xi_j \ge c_\circ\,|\xi|^2\,, \quad c_\circ > 0\,, \tag{3.7}$$

we define one-sided, first-order, trace operators

$$\gamma_1^\pm \in \mathsf{B}(H^s(\Omega_\pm), H^{s-\frac{3}{2}}(\Gamma))\,, \qquad \frac{3}{2} < s \le k+1\,, \tag{3.8}$$

by the zero-order trace of the co-normal derivative:

$$\gamma_1^\pm u := \sum_{1\le i,j\le n} \nu_i\gamma_0^\pm(a_{ij}\partial_{x_j}u) \tag{3.9}$$

Using the maps $\gamma_0^\pm$ and $\gamma_1^\pm$ we define the two-sided, bounded, trace operators

$$\gamma_0 : H^s(\Omega_-)\oplus H^s(\Omega_+) \to H^{s-\frac{1}{2}}(\Gamma)\,, \quad \gamma_0(u_-\oplus u_+) := \frac{1}{2}(\gamma_0^+u_+ + \gamma_0^-u_-)\,,$$

$$\gamma_1 : H^s(\Omega_-)\oplus H^s(\Omega_+) \to H^{s-\frac{3}{2}}(\Gamma)\,, \quad \gamma_1(u_-\oplus u_+) := \frac{1}{2}(\gamma_1^+u_+ + \gamma_1^-u_-)$$

and

$$[\gamma_0] : H^s(\Omega_-)\oplus H^s(\Omega_+) \to H^{s-\frac{1}{2}}(\Gamma)\,, \quad [\gamma_0](u_-\oplus u_+) := \gamma_0^+u_+ - \gamma_0^-u_-\,,$$

$$[\gamma_1] : H^s(\Omega_-) \oplus H^s(\Omega_+) \to H^{s-\frac{3}{2}}(\Gamma)\,, \quad [\gamma_1](u_- \oplus u_+) := \gamma_1^+ u_+ - \gamma_1^- u_-\,.$$

Posing

$$H^s(\mathbb{R}^n\backslash\Gamma) := H^s(\Omega_-) \oplus H^s(\Omega_+)\,,$$

by [1, Theorem 3.5.1], one has

$$H^s(\mathbb{R}^n) = H^s(\mathbb{R}^n\backslash\Gamma)\,, \quad 0 \le s < \frac{1}{2}\,, \tag{3.10}$$

$$H^s(\mathbb{R}^n) = H^s(\mathbb{R}^n\backslash\Gamma) \cap \ker([\gamma_0])\,, \quad \frac{1}{2} < s < \frac{3}{2}\,, \tag{3.11}$$

$$H^s(\mathbb{R}^n) = H^s(\mathbb{R}^n\backslash\Gamma) \cap \ker([\gamma_0]) \cap \ker([\gamma_1]) \quad \frac{3}{2} < s < \frac{5}{2}\,. \tag{3.12}$$

More generally, given a relatively open subset $\Sigma \subset \Gamma$, one has

$$H^s(\mathbb{R}^n\backslash\overline{\Sigma}) = \{u \in H^s(\mathbb{R}^n\backslash\Gamma) : \operatorname{supp}([\gamma_0]u) \subseteq \overline{\Sigma}\}\,, \quad \frac{1}{2} < s < \frac{3}{2}\,, \tag{3.13}$$

and

$$H^s(\mathbb{R}^n\backslash\overline{\Sigma}) = \{u \in H^s(\mathbb{R}^n\backslash\Gamma) : \operatorname{supp}([\gamma_0]u) \cup \operatorname{supp}([\gamma_1]u) \subseteq \overline{\Sigma}\}\,, \quad \frac{3}{2} < s < \frac{5}{2}\,. \tag{3.14}$$

In the following we use the notations

$$H^{s-}(\mathbb{R}^n) := \bigcap_{r<s} H^r(\mathbb{R}^n)\,, \quad H^{s-}(\mathbb{R}^n\backslash\Gamma) := \bigcap_{r<s} H^r(\mathbb{R}^n\backslash\Gamma)\,, \quad H^{s-}(\mathbb{R}^n\backslash\overline{\Sigma}) := \bigcap_{r<s} H^r(\mathbb{R}^n\backslash\overline{\Sigma})\,.$$

3.4. *Boundary-layer operators.*

In what follows A denotes the 2nd order, symmetric, elliptic partial differential operator

$$A : \mathscr{D}'(\mathbb{R}^n) \to \mathscr{D}'(\mathbb{R}^n)\,, \quad Au := \sum_{1\le i,j\le n} \partial_{x_i}(a_{ij}\partial_{x_j}u) - Vu\,,$$

where we suppose a_{ij}, $\partial_{x_i}a_{ij}$, $V \in C_b^\infty(\mathbb{R}^n)$, $a_{ij}(x) = a_{ji}(x)$ and that (3.7) holds true. When restricted to $H^1(\mathbb{R}^n)$, A provides a bounded operator in $\mathsf{B}(H^1(\mathbb{R}^n), H^{-1}(\mathbb{R}^n))$ by the identity

$$\langle Au, v\rangle_{-1,1} = -\sum_{1\le i,j\le n} \langle a_{ij}\partial_{x_i}u, \partial_{x_j}v\rangle_{L^2(\mathbb{R}^n)} + \langle Vu, v\rangle_{L^2(\mathbb{R}^n)}\,.$$

Moreover the sesquilinear form

$$F : H^1(\mathbb{R}^n) \times H^1(\mathbb{R}^n) \subset L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n) \to \mathbb{R}\,, \quad F(u,v) := -\langle Au, v\rangle_{-1,1}$$

satisfies

$$\forall u \in H^1(\mathbb{R}^n)\,, \quad F(u,u) \ge c_\circ \|\nabla u\|^2_{L^2(\mathbb{R}^n)} - \|V_{\text{neg}}\|_\infty \|u\|^2_{L^2(\mathbb{R}^n)}\,, \tag{3.15}$$

where V_{neg} denotes the negative part of V; therefore F is closed and semibounded . By (7.1), the corresponding self-adjoint operator is then given by the restriction of A to the domain D_A,

$$D_A := \{u \in H^1(\mathbb{R}^n) : Au \in L^2(\mathbb{R}^n)\} \equiv H^2(\mathbb{R}^n)\,.$$

Thus $A : H^2(\mathbb{R}^n) \subset L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ is self-adjoint, $(-A+z)^{-1} \in \mathsf{B}(L^2(\mathbb{R}^n), H^2(\mathbb{R}^n))$ for any $z \in \rho(A)$ and $(\|V_{\text{neg}}\|_\infty, +\infty) \subseteq \rho(A)$. By (3.15), for any $\lambda > \|V_{\text{neg}}\|_\infty$ one obtains

$$\forall u \in H^1(\mathbb{R}^n)\,, \quad \|(-A+\lambda)u\|_{H^{-1}(\mathbb{R}^n)} \ge c\,\|u\|_{H^1(\mathbb{R}^n)}\,, \tag{3.16}$$

and so

$$(-A+\lambda)^{-1} \in \mathsf{B}(H^{-1}(\mathbb{R}^n), H^1(\mathbb{R}^n))\,. \tag{3.17}$$

By (3.17) and by elliptic regularity, see e.g. [40, Theorem 6.22], $(-A+\lambda) \in \mathsf{B}(H^{m+2}(\mathbb{R}^n), H^m(\mathbb{R}^n))$ is a bijection; thus, by the inverse mapping theorem, one has

$$(-A+\lambda)^{-1} \in \mathsf{B}(H^m(\mathbb{R}^n), H^{m+2}(\mathbb{R}^n)) \,, \quad m \geq -1 \,. \tag{3.18}$$

Given the bounded open set $\Omega \subset \mathbb{R}^n$ of class $C^{k,1}$, $k \geq 0$, the single and double-layer operators

$$\mathrm{SL}_z : H^{-\frac{3}{2}}(\Gamma) \to L^2(\mathbb{R}^n) \,, \qquad \mathrm{DL}_z : H^{-\frac{1}{2}}(\Gamma) \to L^2(\mathbb{R}^n) \,,$$

related to A and Γ are defined by

$$\langle \mathrm{SL}_z\phi, u\rangle_{L^2(\mathbb{R}^n)} := \langle \phi, \gamma_0(-A+\bar{z})^{-1}u\rangle_{-\frac{3}{2},\frac{3}{2}} \,, \quad u \in L^2(\mathbb{R}^n) \,, \tag{3.19}$$

$$\langle \mathrm{DL}_z\varphi, u\rangle_{L^2(\mathbb{R}^n)} := \langle \varphi, \gamma_1(-A+\bar{z})^{-1}u\rangle_{-\frac{1}{2},\frac{1}{2}} \,, \quad u \in L^2(\mathbb{R}^n) \,. \tag{3.20}$$

Let $g_z(x,y)$ be the integral kernel of the resolvent $(-A+z)^{-1}$; it is a smooth function for $x \neq y$ (see e.g. [57, Lemma 6.3]). Therefore (3.19) and (3.20) give, if $x \notin \Gamma$ and ϕ, $\varphi \in L^2(\Gamma)$,

$$\mathrm{SL}_z\phi(x) = \int_\Gamma g_z(x,y)\,\phi(y)\,d\sigma_\Gamma(y) \,, \tag{3.21}$$

and

$$\mathrm{DL}_z\varphi(x) = \sum_{1\leq i,j\leq n} \int_\Gamma \nu_i(y)a_{ij}(y)\partial_{x_j}g_z(x,y)\,\varphi(y)\,d\sigma_\Gamma(y) \,, \tag{3.22}$$

where σ_Γ denotes the surface measure. We need the following mapping properties:

Lemma 3.1. *For any $\lambda > \|V_{neg}\|_\infty$ one has*

$$\mathrm{SL}_\lambda \in \mathsf{B}(H^{-\frac{1}{2}}(\Gamma), H^1(\mathbb{R}^n)) \,, \qquad \mathrm{DL}_\lambda \in \mathsf{B}(H^{\frac{1}{2}}(\Gamma), H^1(\Omega_\pm))$$

Proof. By (3.17), (3.19) and by $\gamma_0 \in \mathsf{B}(H^1(\mathbb{R}^n), H^{\frac{1}{2}}(\Gamma))$, if $\phi \in H^{-\frac{1}{2}}(\Gamma)$ then

$$|\langle \mathrm{SL}_\lambda\phi, u\rangle_{L^2(\mathbb{R}^n)}| \leq \|\phi\|_{H^{-\frac{1}{2}}(\Gamma)}\|\gamma_0(-A+\lambda)^{-1}u\|_{H^{\frac{1}{2}}(\Gamma)} \leq c\,\|\phi\|_{H^{-\frac{1}{2}}(\Gamma)}\|u\|_{H^{-1}(\mathbb{R}^n)} \,.$$

By [57, Lemma 4.3] and [57, Theorem 3.30] one has

$$\|\gamma_1(-A+\lambda)^{-1}u\|_{H^{-\frac{1}{2}}(\Gamma)} \leq c\left(\|(-A+\lambda)^{-1}u\|_{H^1(\Omega_\pm)} + \|u\|_{H^1(\Omega_\pm)'}\right) \,.$$

Thus, by (3.17) and (3.20), if $\varphi \in H^{\frac{1}{2}}(\Gamma)$ there follows

$$|\langle \mathrm{DL}_\lambda\varphi, u\rangle_{L^2(\Omega_\pm)}| \leq \|\varphi\|_{H^{\frac{1}{2}}(\Gamma)}\|\gamma_1(-A+\lambda)^{-1}1_{\Omega_\pm}u\|_{H^{-\frac{1}{2}}(\Gamma)} \leq c\,\|\varphi\|_{H^{\frac{1}{2}}(\Gamma)}\|u\|_{H^1(\Omega_\pm)'} \,.$$

□

Since $g_z(x,y)$ is a smooth function for $x \neq y$, one has $\mathrm{SL}_z\phi$, $\mathrm{DL}_z\varphi \in C^\infty(\mathbb{R}^n\backslash\Gamma)$ and (see [57, eqs. (6.18) and (6.19)])

$$\forall x \notin \Gamma \,, \qquad A\,\mathrm{SL}_z\phi(x) = z\,\mathrm{SL}_z\phi(x) \,, \quad A\,\mathrm{DL}_z\varphi(x) = z\,\mathrm{DL}_z\varphi(x) \,. \tag{3.23}$$

Therefore, setting

$$\mathrm{SL}_z^\pm\phi := \mathrm{SL}_z\phi|\Omega_\pm \,, \quad \mathrm{DL}_z^\pm\varphi := \mathrm{DL}_z\varphi|\Omega_\pm \,,$$

one has

$$\mathrm{SL}_z^\pm\phi \in \ker(A_\pm^{\max} - z) \,, \qquad \mathrm{DL}_z^\pm\varphi \in \ker(A_\pm^{\max} - z) \,, \tag{3.24}$$

where

$$A_\pm^{\max} = A|\mathrm{dom}(A_\pm^{\max}) \,, \quad \mathrm{dom}(A_\pm^{\max}) := \{u_\pm \in L^2(\Omega_\pm) : Au_\pm \in L^2(\Omega_\pm)\} \,. \tag{3.25}$$

In the case Ω is of class $C^{1,1}$, by proceeding as in the proof of theorem 6.5 in [58, Section 6, Chapter 2] (see the comment in [39] before Theorem 1.5.3.4), the maps $\gamma_0^\pm$ and $\gamma_1^\pm$ can be extended to

$$\hat{\gamma}_0^\pm \in \mathsf{B}(\mathrm{dom}(A_\pm^{\max}), H^{-\frac{1}{2}}(\Gamma)) \,, \quad \hat{\gamma}_1^\pm \in \mathsf{B}(\mathrm{dom}(A_\pm^{\max}), H^{-\frac{3}{2}}(\Gamma))$$

(here $\mathrm{dom}(A^{\pm}_{\max})$ has the graph norm), which in turn provide us with the bounded maps

$$\hat{\gamma}_0 : \mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+}) \to H^{-\frac{1}{2}}(\Gamma)\,, \quad \hat{\gamma}_0(u_- \oplus u_+) := \frac{1}{2}(\hat{\gamma}^+_0 u_+ + \hat{\gamma}^-_0 u_-)\,,$$

$$\hat{\gamma}_1 : \mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+}) \to H^{-\frac{3}{2}}(\Gamma)\,, \quad \hat{\gamma}_1(u_- \oplus u_+) := \frac{1}{2}(\hat{\gamma}^+_1 u_+ + \hat{\gamma}^-_1 u_-)$$

and

$$[\hat{\gamma}_0] : \mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+}) \to H^{-\frac{1}{2}}(\Gamma)\,, \quad [\hat{\gamma}_0](u_- \oplus u_+) := \hat{\gamma}^+_0 u_+ - \hat{\gamma}^-_0 u_-\,,$$

$$[\hat{\gamma}_1] : \mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+}) \to H^{-\frac{3}{2}}(\Gamma)\,, \quad [\hat{\gamma}_1](u_- \oplus u_+) := \hat{\gamma}^+_1 u_+ - \hat{\gamma}^-_1 u_-\,.$$

These maps, together with [57, Theorem 7.2], give, whenever Ω is of class $C^{k,1}$, $k \geq 1$, and for any $|s| \leq k$, the bounded operators

$$\hat{\gamma}^{\pm}_0 \mathrm{SL}_z \in \mathsf{B}(H^{s-\frac{1}{2}}(\Gamma), H^{s+\frac{1}{2}}(\Gamma))\,, \qquad \hat{\gamma}^{\pm}_1 \mathrm{SL}_z \in \mathsf{B}(H^{s-\frac{1}{2}}(\Gamma), H^{s-\frac{1}{2}}(\Gamma))\,, \tag{3.26}$$

$$\hat{\gamma}^{\pm}_0 \mathrm{DL}_z \in \mathsf{B}(H^{s+\frac{1}{2}}(\Gamma), H^{s+\frac{1}{2}}(\Gamma))\,, \qquad \hat{\gamma}^{\pm}_1 \mathrm{DL}_z \in \mathsf{B}(H^{s+\frac{1}{2}}(\Gamma), H^{s-\frac{1}{2}}(\Gamma))\,. \tag{3.27}$$

Moreover the single and double layer operators satisfy the jump relations

$$[\hat{\gamma}_0]\mathrm{SL}_z\phi = [\hat{\gamma}_1]\mathrm{DL}_z\phi = 0\,, \quad [\hat{\gamma}_1]\mathrm{SL}_z\phi = -[\hat{\gamma}_0]\mathrm{DL}_z\phi = -\phi\,. \tag{3.28}$$

By the first relation in (3.28) one gets

$$\hat{\gamma}^{\pm}_0 \mathrm{SL}_z = \hat{\gamma}_0 \mathrm{SL}_z\,, \qquad \hat{\gamma}^{\pm}_1 \mathrm{DL}_z = \hat{\gamma}_1 \mathrm{DL}_z\,.$$

Notice that in (3.26), (3.27) and (3.28) the extended trace operators coincide with the usual ones whenever the range spaces are Sobolev space on Γ of strictly positive index.

For any $\lambda \in \rho(A) \cap \mathbb{R}$ both the bounded operators $\gamma_0\mathrm{SL}_\lambda$ and $\hat{\gamma}_1\mathrm{DL}_\lambda$ are symmetric w.r.t. the $H^{-\frac{1}{2}}(\Gamma)$-$H^{\frac{1}{2}}(\Gamma)$ pairing (see [57, Theorems 6.15 and 6.17]:

$$\forall \phi, \varphi \in H^{-\frac{1}{2}}(\Gamma)\,, \quad \langle \phi, \gamma_0\mathrm{SL}_\lambda\varphi\rangle_{-\frac{1}{2},\frac{1}{2}} = \langle \gamma_0\mathrm{SL}_\lambda\phi, \varphi\rangle_{\frac{1}{2},-\frac{1}{2}}\,,$$

$$\forall \phi, \varphi \in H^{\frac{1}{2}}(\Gamma)\,, \quad \langle \phi, \hat{\gamma}_1\mathrm{DL}_\lambda\varphi\rangle_{\frac{1}{2},-\frac{1}{2}} = \langle \hat{\gamma}_1\mathrm{DL}_\lambda\phi, \varphi\rangle_{-\frac{1}{2},\frac{1}{2}}\,.$$

Moreover these operators are coercive:

Lemma 3.2. *Let $\lambda > \max(V_{neg})$. Then there exist $c_0 > 0$ and $c_1 > 0$ such that*

$$\forall \phi \in H^{-\frac{1}{2}}(\Gamma)\,, \quad \langle \phi, \gamma_0\mathrm{SL}_\lambda\phi\rangle_{-\frac{1}{2},\frac{1}{2}} \geq c_0\, \|\phi\|^2_{H^{-\frac{1}{2}}(\Gamma)} \tag{3.29}$$

and

$$\forall \varphi \in H^{\frac{1}{2}}(\Gamma)\,, \quad -\langle \hat{\gamma}_1\mathrm{DL}_\lambda\varphi, \varphi\rangle_{-\frac{1}{2},\frac{1}{2}} \geq c_1\, \|\varphi\|^2_{H^{\frac{1}{2}}(\Gamma)}\,. \tag{3.30}$$

Proof. In the case $A = \Delta$ and $\lambda = 1$, the proof is given in [50, Lemma 1.14 (c)] as regards $\hat{\gamma}_0\mathrm{SL}_\lambda$ and in [50, Theorem 1.26 (e)] as regards $\hat{\gamma}_1\mathrm{DL}_\lambda$. Here we provide an alternative proof which adapts to our hypotheses.

By [57, Lemma 4.3] for any $u_\pm \in \mathrm{dom}(A^{\max}_{\pm}) \cap H^1(\Omega_\pm)$ one has $\hat{\gamma}^{\pm}_1 u_\pm \in H^{-\frac{1}{2}}(\Gamma)$ and

$$\|\hat{\gamma}^{\pm}_1 u_\pm\|^2_{H^{-\frac{1}{2}}(\Gamma)} \leq c\left(\|u_\pm\|^2_{H^1(\Omega_\pm)} + \|A^{\max}_{\pm} u_\pm\|^2_{H^{-1}(\Omega_\pm)}\right)\,; \tag{3.31}$$

moreover for such a $u_\pm$ and for any $v_\pm \in H^1(\Omega_\pm)$ the ”half” Green's formula holds (see [57, Theorem 4.4]):

$$\begin{aligned}&\langle(-A^{\max}_{\pm} + \lambda)u_\pm, v_\pm\rangle_{L^2(\Omega_\pm)}\\ &= \sum_{1\leq i,j\leq n} \langle a_{ij}\partial_i u_\pm, \partial_j v_\pm\rangle_{L^2(\Omega_\pm)} + \langle(V+\lambda)u_\pm, v_\pm\rangle_{L^2(\Omega_\pm)} \pm \langle \hat{\gamma}^{\pm}_1 u_\pm, \gamma^{\pm}_0 v_\pm\rangle_{-\frac{1}{2},\frac{1}{2}}\,.\end{aligned}$$

Thus, posing $u_\pm = v_\pm = \mathrm{SL}^{\pm}_{\lambda}\phi$, by $(-A^{\pm}_{\max} + \lambda)u_\pm = 0$ and by (3.28), one gets

$$\begin{aligned}0 = &\sum_{1\leq i,j\leq n} \langle a_{ij}\partial_i\mathrm{SL}_\lambda\phi, \partial_j\mathrm{SL}_\lambda\phi\rangle_{L^2(\mathbb{R}^n)} + \langle(V+\lambda)\mathrm{SL}_\lambda\phi, \mathrm{SL}_\lambda\phi\rangle_{L^2(\mathbb{R}^n)} + \langle[\hat{\gamma}_1]\mathrm{SL}_\lambda\phi, \gamma_0\mathrm{SL}_\lambda\phi\rangle_{-\frac{1}{2},\frac{1}{2}}\\ \geq &c_\circ\|\nabla\mathrm{SL}_\lambda\phi\|^2_{L^2(\mathbb{R}^n)} + (\lambda - \max(V_{\mathrm{neg}}))\,\|\mathrm{SL}_\lambda\phi\|^2_{L^2(\mathbb{R}^n)} - \langle\phi, \gamma_0\mathrm{SL}_z\phi\rangle_{-\frac{1}{2},\frac{1}{2}}\,.\end{aligned}$$

Therefore, setting $\kappa_\circ := \min\{c_\circ, \lambda - \max(V_{\text{neg}})\}$, one obtains

$$\langle \phi, \gamma_0 \mathrm{SL}_\lambda \phi\rangle_{-\frac{1}{2},\frac{1}{2}} \geq \kappa_\circ \, \|\mathrm{SL}_\lambda \phi\|^2_{H^1(\mathbb{R}^n)} .$$

By $[\hat{\gamma}_1]\mathrm{SL}_\lambda\phi = -\phi$, by (3.31) and by the continuous embedding $H^1(\Omega_\pm) \hookrightarrow H^{-1}(\Omega_\pm)$ one gets

$$\begin{aligned}\|\phi\|^2_{H^{-\frac{1}{2}}(\Gamma)} \leq & c\left(\|\mathrm{SL}^-_\lambda\phi\|^2_{H^1(\Omega_-)} + \lambda^2\|\mathrm{SL}^-_\lambda\phi\|^2_{H^{-1}(\Omega_-)} + \|\mathrm{SL}^+_\lambda\phi\|^2_{H^1(\Omega_+)} + \lambda^2\|\mathrm{SL}^+_\lambda\phi\|^2_{H^{-1}(\Omega)}\right) \\ \leq & c\,(1+\lambda^2)\|\mathrm{SL}_\lambda\phi\|^2_{H^1(\mathbb{R}^n)}\end{aligned}$$

and so (3.29) follows by posing $c_0 = \kappa_\circ(c\,(1+\lambda^2))^{-1}$.

The proof of (3.30) proceeds along the same lines by inserting $u_\pm = v_\pm = \mathrm{DL}^\pm_\lambda\varphi$ in the Green's formula above: in this case one obtains

$$-\langle \hat{\gamma}_1 \mathrm{DL}_\lambda\varphi, \varphi\rangle_{-\frac{1}{2},\frac{1}{2}} \geq \kappa_0 \, \|\mathrm{DL}_\lambda\varphi\|^2_{H^1(\Omega_-)\oplus H^1(\Omega_+)} .$$

By $[\gamma_0]\mathrm{DL}_\lambda\varphi = \varphi$, denoting by $N_\pm$ the norm of $\gamma_0^\pm \in \mathsf{B}(H^1(\Omega_\pm), H^{\frac{1}{2}}(\Gamma))$, one obtains

$$\|\varphi\|^2_{H^{\frac{1}{2}}(\Gamma)} \leq N^2_-\|\mathrm{DL}^-_\lambda\varphi\|^2_{H^1(\Omega_-)} + N^2_+\|\mathrm{DL}^+_\lambda\varphi\|^2_{H^1(\Omega_+)} \leq \max\{N^2_-, N^2_+\}\|\mathrm{DL}_\lambda\varphi\|^2_{H^1(\Omega_-)\oplus H^1(\Omega_+)} ,$$

and so (3.29) follows by posing $c_1 = \kappa_\circ(\max\{N^2_-, N^2_+\})^{-1}$. ☐

We conclude this section providing results about the mapping properties of $\hat{\gamma}_0\mathrm{SL}_\lambda$ and $\hat{\gamma}_1\mathrm{DL}_\lambda$.

Lemma 3.3. *Let $\lambda > \max(V_{neg})$ and Ω be of class $C^{k,1}$, $k \geq 1$. Then*

$$\hat{\gamma}_0\mathrm{SL}_\lambda\phi \in H^{s+\frac{1}{2}}(\Gamma) \quad\iff\quad \phi \in H^{s-\frac{1}{2}}(\Gamma)\,, \quad |s| \leq k-1\,, \tag{3.32}$$

$$\hat{\gamma}_1\mathrm{DL}_\lambda\phi \in H^{s-\frac{1}{2}}(\Gamma) \quad\iff\quad \phi \in H^{s+\frac{1}{2}}(\Gamma)\,, \qquad |s| \leq k\,. \tag{3.33}$$

Proof. By (3.26) and (3.27) we only need to prove the $\Rightarrow$ implications. By [57, Theorem 7.17], both the maps $\hat{\gamma}_0\mathrm{SL}_\lambda$ and $\hat{\gamma}_1\mathrm{DL}_\lambda$ are Fredholm with index zero and s-independent kernel. By (3.29) and (3.30) such maps are injective and therefore bijective. Thus, by (3.26), (3.27) and by the inverse mapping theorem, $(\hat{\gamma}_0\mathrm{SL}_\lambda)^{-1} \in \mathsf{B}(H^{s+\frac{1}{2}}(\Gamma), H^{s-\frac{1}{2}}(\Gamma))$ and $(\hat{\gamma}_1\mathrm{DL}_\lambda)^{-1} \in \mathsf{B}(H^{s-\frac{1}{2}}(\Gamma), H^{s+\frac{1}{2}}(\Gamma))$. ☐

4. Self-adjoint realizations of singular perturbations supported on hypersurfaces.

Let $A : H^2(\mathbb{R}^n) \subset L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ be the elliptic, self-adjoint operator defined in Subsection 3.4, i.e.

$$Au := \sum_{1\leq i,j\leq n} \partial_{x_i}(a_{ij}\partial_{x_j}u) - Vu\,,$$

with $a_{ij}, \partial_{x_i}a_{ij}, V \in C^\infty_b(\mathbb{R}^n)$, the symmetric matrix $\underline{a} \equiv [a_{ij}(x)]$ satisfying (3.7).

Given Ω open, bounded and of class $C^{1,1}$, posing

$$\tau : H^2(\mathbb{R}^n) \to H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)\,, \quad \tau u := \gamma_0 u \oplus \gamma_1 u\,,$$

one has

Lemma 4.1. *The map τ is bounded, surjective and* $\ker(\tau)$ *is dense in* $L^2(\mathbb{R}^n)$.

Proof. Let $u \in H^2(\mathbb{R}^n)$. Then, by $\|u\|^2_{H^2(\mathbb{R}^n)} = \|u|\Omega_-\|^2_{H^2(\Omega_-)} + \|u|\Omega_+\|^2_{H^2(\Omega_+)}$, τ is bounded since both $\gamma_0^\pm$ and $\gamma_1^\pm$ are bounded; $\ker(\tau)$ is dense since it contains the dense set $C^\infty_{\text{comp}}(\mathbb{R}^n\backslash\Gamma)$. By [39, Remark 2.5.1.2], both

$$\tau^\pm : H^2(\Omega_\pm) \to H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)\,, \quad \tau^\pm u_\pm := \gamma_0^\pm u_\pm \oplus \gamma_1^\pm u_\pm$$

are surjective. Given $\phi \oplus \varphi \in H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)$, let $u_\pm \in H^2(\Omega_\pm)$ such that $\tau^\pm u_\pm = \phi \oplus \varphi$. Let $\tilde{u}_\pm \in H^2(\mathbb{R}^n)$ be an extensions of $u_\pm$. Then $\tau\left(\frac{1}{2}\,(\tilde{u}_- + \tilde{u}_+)\right) = \phi \oplus \varphi$ and so τ is surjective. ☐

By the definition of the map τ we have that

$$S := A|\ker(\tau) = A_-^{\min} \oplus A_+^{\min}\,,$$

where

$$A_\pm^{\min} := A|H_0^2(\Omega_\pm)\,, \quad H_0^2(\Omega_\pm) := \{u_\pm \in H^2(\Omega_\pm) : \gamma_0^\pm u_\pm = \gamma_1^\pm u_\pm = 0\}\,.$$

Since it is known that $(A_\pm^{\min})^* = A_\pm^{\max}$, one obtains

$$\mathrm{dom}(S^*) = \mathrm{dom}(A_-^{\max}) \oplus \mathrm{dom}(A_+^{\max})\,, \qquad S^* = A_-^{\max} \oplus A_+^{\max}\,. \tag{4.1}$$

Thus we have the well-defined bounded (w.r.t. the graph norm in $\mathrm{dom}(S^*)$) maps

$$\hat\gamma_0,\, [\hat\gamma_0] \in \mathsf{B}(\mathrm{dom}(S^*), H^{-\frac{1}{2}}(\Gamma))\,, \quad \hat\gamma_1,\, [\hat\gamma_1] \in \mathsf{B}(\mathrm{dom}(S^*), H^{-\frac{3}{2}}(\Gamma))\,.$$

By Lemma 4.1 we can apply the results of Section 2 to A and so find all self-adjoint extensions of the closed symmetric operator $A|\ker(\tau) = A_-^{\min} \oplus A_+^{\min}$. To this end we need to determine the operator G_z and M_z (see definitions (2.1) and (2.6)). By (3.19) and (3.20) this is immediate:

$$G_z : H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma) \to L^2(\mathbb{R}^n)\,, \quad G_z(\phi \oplus \varphi) := \mathrm{SL}_z\phi + \mathrm{DL}_z\varphi \tag{4.2}$$

and

$$\begin{gathered} M_z : H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)\,, \\ M_z := \begin{bmatrix} \gamma_0(\mathrm{SL} - \mathrm{SL}_z) & \gamma_0(\mathrm{DL} - \mathrm{DL}_z) \\ \gamma_1(\mathrm{SL} - \mathrm{SL}_z) & \gamma_1(\mathrm{DL} - \mathrm{DL}_z) \end{bmatrix}, \end{gathered} \tag{4.3}$$

where

$$\mathrm{SL} := \mathrm{SL}_{\lambda_\circ}\,, \qquad \mathrm{DL} := \mathrm{DL}_{\lambda_\circ}\,, \qquad \lambda_\circ > \max(V_{\mathrm{neg}})\,.$$

Next lemma provides a representation of $A_-^{\max} \oplus A_+^{\max}$ and of its domain. Before giving the precise statement we need some definitions. The distribution $\delta_\Gamma \in \mathscr{D}'(\mathbb{R}^n)$ is defined as usual by

$$\forall u \in C^\infty_{\mathrm{comp}}(\mathbb{R}^n)\,, \quad (\delta_\Gamma, u) = \int_\Gamma u(x)\, d\sigma_\Gamma(x)\,.$$

Given $f \in H^{-s}(\Gamma)$, we then define $f\delta_\Gamma \in \mathscr{D}'(\mathbb{R}^n)$ and $f\partial_{\underline{a}}\delta_\Gamma \in \mathscr{D}'(\mathbb{R}^n)$ by

$$\forall u \in C^\infty_{\mathrm{comp}}(\mathbb{R}^n)\,, \quad (f\delta_\Gamma, u) = \langle \bar{f}, u|\Gamma\rangle_{-s,s}$$

and

$$\forall u \in C^\infty_{\mathrm{comp}}(\mathbb{R}^n)\,, \quad (f\partial_{\underline{a}}\delta_\Gamma, u) := -\sum_{1\le i,j\le n} (f\nu_i\delta_\Gamma, a_{ij}\partial_{x_j}u)\,.$$

Notice that if Ω is of class $C^{1,1}$ then ν is Lipschitz continuous and so the product $f\nu$ is a well-defined vector in $H^{-r}(\Gamma)$, $r = \min\{1, s\}$.

Lemma 4.2.

$$\begin{aligned} \mathrm{dom}(A_-^{\max}) \oplus \mathrm{dom}(A_+^{\max}) =& \{u = u_\circ + \mathrm{SL}\phi + \mathrm{DL}\varphi\,,\ u_\circ \in H^2(\mathbb{R}^n)\,,\ \phi \oplus \varphi \in H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma)\} \\ \equiv& \{u = u_\circ - \mathrm{SL}[\hat\gamma_1]u + \mathrm{DL}[\hat\gamma_0]u\,,\ u_\circ \in H^2(\mathbb{R}^n)\}\,, \end{aligned}$$

and

$$(A_-^{\max} \oplus A_+^{\max})u = Au - [\hat\gamma_1]u\,\delta_\Gamma - [\hat\gamma_0]u\,\partial_{\underline{a}}\delta_\Gamma\,. \tag{4.4}$$

Proof. By (4.1), Lemma 2.3 and (4.2) one has

$$\mathrm{dom}(A_-^{\max}) \oplus \mathrm{dom}(A_+^{\max}) = \{u = u_\circ + G(\phi \oplus \varphi)\,,\ u_\circ \in H^2(\mathbb{R}^n)\,,\ \phi \oplus \varphi \in H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma)\}\,.$$

Since $[\gamma_0]u_\circ = [\gamma_1]u_\circ = 0$, the proof of the first statement follows by using the jump relations (3.28). As regards the second statement, in the case $A = \Delta$ the proof has been given in [21, Theorem 3.1] (be aware that there the jumps of the trace maps have been defined with opposite signs). The proof in the more general case discussed here proceeds along the same lines and is left to the reader. ☐

Remark 4.3. By $(A_-^{\max} \oplus A_+^{\max})G_z(\phi \oplus \varphi) = zG_z(\phi \oplus \varphi)$ (see (2.8)) and by (4.4), one has $AG_z(\phi \oplus \varphi)(x) = zG_z(\phi \oplus \varphi)(x)$ for any $x \in \mathbb{R}^n \backslash \Gamma$. Hence, by elliptic regularity (see e.g. [57, Theorem 6.4]), setting $\Gamma^\epsilon := \{x \in \mathbb{R}^n : \operatorname{dist}(x, \Gamma) \leq \epsilon\}$, $\epsilon > 0$, one has

$$\forall (\phi \oplus \varphi) \in H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma) , \qquad G_z(\phi \oplus \varphi) \in \bigcap_{s>0} H^s(\mathbb{R}^n \backslash \Gamma^\epsilon) . \tag{4.5}$$

Similarly, if $\operatorname{supp}(\phi) \subseteq \overline{\Sigma}$ and $\operatorname{supp}(\varphi) \subseteq \overline{\Sigma}$, $\Sigma \subset \Gamma$ relatively open, then by (4.4), one has $AG_z(\phi \oplus \varphi)(x) = zG_z(\phi \oplus \varphi)(x)$ for any $x \in \mathbb{R}^n \backslash \overline{\Sigma}$. Hence, by [57, Theorem 6.4], setting $\Sigma^\epsilon := \{x \in \mathbb{R}^n : \operatorname{dist}(x, \Sigma) \leq \epsilon\}$, $\epsilon > 0$,

$$\forall (\phi \oplus \varphi) \in H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) \oplus H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) , \qquad G_z(\phi \oplus \varphi) \in \bigcap_{s>0} H^s(\mathbb{R}^n \backslash \Sigma^\epsilon) . \tag{4.6}$$

From now on

$$\Pi : H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma) ,$$

denotes an orthogonal projector,

$$\Pi' : H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma) \to H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma) ,$$

denotes the orthogonal projector defined as the dual of Π, so that $\Pi' = (\Lambda^3 \oplus \Lambda)\Pi(\Lambda^{-3} \oplus \Lambda^{-1})$, and

$$\Theta : \operatorname{dom}(\Theta) \subseteq \operatorname{ran}(\Pi') \to \operatorname{ran}(\Pi)$$

denotes a self-adjoint operator. By Theorem 2.1 and Lemma 4.2, one readily obtains all self-adjoint extension of S:

Theorem 4.4. *Any self-adjoint extension of $A_-^{\min} \oplus A_+^{\min}$ is of the kind $A_{\Pi,\Theta}$, where*

$$A_{\Pi,\Theta} : \operatorname{dom}(A_{\Pi,\Theta}) \subseteq L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n) , \qquad A_{\Pi,\Theta} := (A_-^{\max} \oplus A_+^{\max})|\operatorname{dom}(A_{\Pi,\Theta}) ,$$

$$\begin{aligned}&\operatorname{dom}(A_{\Pi,\Theta}) := \{u \in \operatorname{dom}(A_-^{\max}) \oplus \operatorname{dom}(A_+^{\max}) : (-[\hat{\gamma}_1]u) \oplus [\hat{\gamma}_0]u \in \operatorname{dom}(\Theta) ,\\ &\Pi(\gamma_0(u + \mathrm{SL}[\hat{\gamma}_1]u - \mathrm{DL}[\hat{\gamma}_0]u) \oplus \gamma_1(u + \mathrm{SL}[\hat{\gamma}_1]u - \mathrm{DL}[\hat{\gamma}_0]u)) = \Theta((-[\hat{\gamma}_1]u) \oplus [\hat{\gamma}_0]u)\} .\end{aligned}$$

The set

$$Z_{\Pi,\Theta} := \{z \in \rho(A) : \Theta + \Pi M_z \Pi' \textit{ has a bounded inverse}\}$$

is not void; in particular $\mathbb{C} \backslash \mathbb{R} \subseteq Z_{\Pi,\Theta} \subseteq \rho(A_{\Pi,\Theta})$ and for any $z \in Z_{\Pi,\Theta}$ the resolvent of $A_{\Pi,\Theta}$ is given by

$$(-A_{\Pi,\Theta} + z)^{-1}u = (-A + z)^{-1}u + G_z\Pi'(\Theta + \Pi M_z \Pi')^{-1}\Pi(\gamma_0((-A + z)^{-1}u) \oplus \gamma_1((-A + z)^{-1}u)) , \tag{4.7}$$

where G_z and M_z are defined in (4.2) *and* (4.3) *respectively.*

Remark 4.5. Let us notice that the Π''s appearing in formula (4.7) act there as the inclusion map $\Pi' : \operatorname{ran}(\Pi') \to H^{-3/2}(\Gamma) \oplus H^{-1/2}(\Gamma)$. This means that one does not need to know Π' explicitly: it suffices to know the subspace $\operatorname{ran}(\Pi') = \operatorname{ran}(\Pi)'$.

Remark 4.6. Let us notice that the self-adjoint extension A corresponds to the choice $\Pi = 0$. By Lemma 4.2, the choice $\Pi = \Pi_1 \oplus 0$ gives $[\hat{\gamma}_0]u = 0$ and so produces self-adjoint extension ("δ-type" interactions) of the kind $Au - [\hat{\gamma}_1]u\,\delta_\Gamma$ while the choice $\Pi = 0 \oplus \Pi_2$ gives $[\hat{\gamma}_1]u = 0$ and so produces self-adjoint extension ("δ'-type" interactions) of the kind $Au - [\hat{\gamma}_0]u\,\partial_{\underline{a}}\delta_\Gamma$; different Π's give combinations of δ and δ' interactions.

Remark 4.7. Let $u = u_\circ + \mathrm{SL}\phi + \mathrm{DL}\varphi$ be in $\operatorname{dom}(A_{\Pi,\Theta})$ and suppose that

$$\operatorname{dom}(\Theta) \subseteq H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma) , \quad s_1 > -\frac{3}{2} , \; s_2 > -\frac{1}{2} . \tag{4.8}$$

Then, by (4.5) and the mapping properties of single and double layer operators (see [57, Corollary 6.14]),

$$\mathrm{SL}\phi + \mathrm{DL}\varphi \in H^s(\Omega_-) \oplus H^s(\Omega_+) , \quad s = \min\left\{s_1 + \frac{3}{2}, s_2 + \frac{1}{2}\right\}$$

and so

$$\operatorname{dom}(A_{\Pi,\Theta}) \subseteq H^{s_\circ}(\mathbb{R}^n \backslash \Gamma) , \quad s_\circ = \min\{2, s\} .$$

In particular, $\operatorname{dom}(A_{\Pi,\Theta}) \subseteq H^2(\mathbb{R}^n \backslash \Gamma)$ whenever $s \geq 2$. The relation (4.6) suggests that such a kind of regularity could hold on a larger set whenever $\operatorname{supp}(\phi) \cup \operatorname{supp}(\varphi) \subseteq \overline{\Sigma}$, $\Sigma \subset \Gamma$. However, as the next result shows, one has to exclude a neighborhood of the interface $\partial\Sigma$:

Lemma 4.8. *Let*

$$D_{1,1} :=\{\phi \in H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) : \phi|\Sigma \in H^{\frac{1}{2}}(\Sigma)\}\,, \qquad D_{2,1} := \{\varphi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \varphi|\Sigma \in H^{\frac{3}{2}}(\Sigma)\}\,,$$
$$D_{1,2} :=\{\phi \in H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) : (\hat\gamma_0\mathrm{SL}\phi)|\Sigma \in H^{\frac{3}{2}}(\Sigma)\}\,, \qquad D_{2,2} := \{\varphi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : (\hat\gamma_1\mathrm{DL}\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma)\}\,.$$

If

$$\mathrm{dom}(\Theta) \subseteq D_{1,i} \oplus D_{2,j}\,, \quad i, j \in \{1, 2\}\,,$$

then

$$\mathrm{dom}(A_{\Pi,\Theta}) \subseteq H^2(\mathbb{R}^n\backslash(\overline{\Sigma} \cup (\partial\Sigma)^\epsilon))\,,$$

where

$$(\partial\Sigma)^\epsilon := \{x \in \mathbb{R}^n : \mathrm{dist}(x, \partial\Sigma) \le \epsilon\}\,, \quad \epsilon > 0\,.$$

Proof. If $\phi \in D_{1,1}$ and $\varphi \in D_{2,1}$, by [57, Theorem 6.13], one has $\mathrm{SL}\phi \in H^2(\Omega^\pm_1)$ and $\mathrm{DL}\varphi \in H^2(\Omega^\pm_1)$, where $\Omega^\pm_1 := \Omega_1 \cap \Omega_\pm$ and Ω_1 of class $C^{1,1}$ such that $\Omega_1 \cap \Gamma$ is strictly contained in Σ.

If $\phi \in D_{1,2}$ and $\varphi \in D_{2,2}$, by [57, Theorem 7.16], $\phi \in H^{\frac{1}{2}}(\Sigma_1)$ and $\varphi \in H^{\frac{3}{2}}(\Sigma_1)$, for any relatively open Σ_1 such that $\overline{\Sigma}_1 \subset \Sigma$. Let now take Ω_2 of class $C^{1,1}$ such that $\Omega_2 \cap \Gamma$ is strictly contained in Σ_1. Then, by [57, Theorem 6.13], one has $\mathrm{SL}\phi \in H^2(\Omega^\pm_2)$ and $\mathrm{DL}\varphi \in H^2(\Omega^\pm_2)$, where $\Omega^\pm_2 := \Omega_2 \cap \Omega_\pm$.

The final statement then follows from (4.6). ☐

In the case $\mathrm{dom}(A_{\Pi,\Theta}) \subseteq H^2(\mathbb{R}^n\backslash\Gamma)$, the statements contained in Theorem 4.4 simplify and one has the following

Corollary 4.9. *Suppose that* $\mathrm{dom}(\Theta) \subseteq H^{\frac{1}{2}}(\Gamma) \oplus H^{\frac{3}{2}}(\Gamma)$ *and define*

$$B_\Theta : \mathrm{dom}(\Theta) \subseteq \mathrm{ran}(\Pi') \to \mathrm{ran}(\Pi)\,, \quad B_\Theta := \Theta + \Pi B\Pi'\,,$$
$$B : H^{\frac{1}{2}}(\Gamma) \oplus H^{\frac{3}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)\,, \quad B := \begin{bmatrix} \gamma_0\mathrm{SL} & \gamma_0\mathrm{DL} \\ \gamma_1\mathrm{SL} & \gamma_1\mathrm{DL} \end{bmatrix}.$$

Then

$$\mathrm{dom}(A_{\Pi,\Theta}) = \{u \in H^2(\mathbb{R}^n\backslash\Gamma) : (-[\gamma_1]u) \oplus [\gamma_0]u \in \mathrm{dom}(\Theta)\,,\ \Pi(\gamma_0u \oplus \gamma_1u) = B_\Theta((-[\gamma_1]u) \oplus [\gamma_0]u)\}$$

and

$$(-A_{\Pi,\Theta} + z)^{-1}u = (-A + z)^{-1}u + G_z\Pi'(B_\Theta - \Pi M^\circ_z\Pi')^{-1}\Pi(\gamma_0(-A + z)^{-1}u \oplus \gamma_1(-A + z)^{-1}u)\,,$$

where

$$M^\circ_z := \begin{bmatrix} \gamma_0\mathrm{SL}_z & \gamma_0\mathrm{DL}_z \\ \gamma_1\mathrm{SL}_z & \gamma_1\mathrm{DL}_z \end{bmatrix}.$$

Proof. By Remark 4.7, $u \in H^2(\Omega_-) \oplus H^2(\Omega_+)$. Thus $\Pi(\gamma_0u \oplus \gamma_1u)$ is well defined and, by the definition of $\mathrm{dom}(A_{\Pi,\Theta})$ given in Theorem 4.4,

$$\begin{aligned}\Theta(\phi \oplus \varphi) &= \Pi(\gamma_0(u - \mathrm{SL}\phi - \mathrm{DL}\varphi) \oplus \gamma_1(u - \mathrm{SL}\phi - \mathrm{DL}\varphi))\\ &=\Pi(\gamma_0u \oplus \gamma_1u) - \Pi(\gamma_0(\mathrm{SL}\phi + \mathrm{DL}\varphi) \oplus \gamma_1(\mathrm{SL}\phi + \mathrm{DL}\varphi))\,.\end{aligned}$$

The proof is then concluded by the identity $\Theta + \Pi M_z\Pi' = B_\Theta - \Pi M^\circ_z\Pi'$. ☐

Let us recall some definitions: $\mathfrak{S}_\infty(H_1, H_2)$ ($\mathfrak{S}_\infty(H) := \mathfrak{S}_\infty(H, H)$), denotes the operator ideal of compact operators on the Hilbert space H_1 to the Hilbert space H_2; $\mathfrak{S}_{p,\infty}(H_1, H_2)$ ($\mathfrak{S}_{p,\infty}(H) := \mathfrak{S}_{p,\infty}(H, H)$), $p > 0$, denote the operator ideals of compact operators T on the Hilbert space H_1 to the Hilbert space H_2 such that $s_k(T) = O(k^{-1/p})$, where the the singular values $s_k(T)$ are defined as the eigenvalues of the non-negative compact operator $(T^*T)^{\frac{1}{2}}$. One has $T_2T_1 \in \mathfrak{S}_{p,\infty}(H_1, H_2)$ whenever $T_1 \in \mathfrak{S}_{p_1,\infty}(H_1, H_0)$, $T_2 \in \mathfrak{S}_{p_2,\infty}(H_0, H_2)$ and $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Notice that if $p < q$ then $\mathfrak{S}_{p,\infty}(H_1, H_2) \subset \mathfrak{S}_q(H_1, H_2)$, where $\mathfrak{S}_q(H_1, H_2)$ denotes the Schatten-von Neumann ideal of compact operator with q-summable singular values; in particular $T \in \mathfrak{S}_{p,\infty}(H_1, H_2)$ is trace class whenever $p < 1$.

Lemma 4.10. *Let* Θ *satisfy* (4.8). *Then*

$$\sigma_{ess}(A_{\Pi,\Theta}) = \sigma_{ess}(A)\,.$$

Proof. Here we follow the same kind of reasonings as in the proof of [11, Lemma 4.7]. The operator $(\Theta + \Pi M_z \Pi')^{-1} \in \mathsf{B}(\mathrm{ran}(\Pi), \mathrm{ran}(\Pi'))$ is closed as an operator from $\mathrm{ran}(\Pi)$ into $H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma)$, hence, since $\mathrm{ran}((\Theta + \Pi M_z \Pi')^{-1}) = \mathrm{dom}(\Theta) \subseteq H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma)$, it is also closed as an operator from $\mathrm{ran}(\Pi)$ into $H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma)$, which implies that it belongs to $\mathsf{B}(\mathrm{ran}(\Pi), H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma))$. Denoting by $E_{s,r}$ the compact embedding of $H^s(\Gamma)$ into $H^r(\Gamma)$, $r < s$, by

$$(\Theta + \Pi M_z \Pi')^{-1} = (E_{s_1,-\frac{3}{2}} \oplus E_{s_2,-\frac{1}{2}})(\Theta + \Pi M_z \Pi')^{-1} ,$$

one gets $(\Theta + \Pi M_z \Pi')^{-1} \in \mathfrak{S}_\infty(\mathrm{ran}(\Pi), \mathrm{ran}(\Pi'))$. Thus, by (4.7), the resolvent difference $(-A_{\Pi,\Theta} + z)^{-1} - (-A + z)^{-1}$ belongs to $\mathfrak{S}_\infty(L^2(\mathbb{R}^n))$ and so $\sigma_{ess}(A_{\Pi,\Theta}) = \sigma_{ess}(A)$ by Weyl's theorem on the preservation of the essential spectrum under compact resolvent perturbations (see e.g. [65, Theorem 8.12]).

□

The next result applies to all the self-adjoint extensions given in Corollary 4.9 (and so to all the operators considered in Section 5). In the proof we follow the same arguments as in [55] and [12].

Theorem 4.11. *Suppose that* Γ *is smooth. If* $\mathrm{dom}(\Theta) \subseteq H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma)$ *with*

$$s = \min\left\{s_1 + \frac{3}{2}, s_2 + \frac{1}{2}\right\} \ge 2 ,$$

then for any integer $k \ge 1$ *and for any* $z \in \rho(A) \cap \rho(A_{\Pi,\Theta})$ *one has*

$$(-A_{\Pi,\Theta} + z)^{-k} - (-A + z)^{-k} \in \mathfrak{S}_{\frac{n-1}{2(k-1)+s},\infty}(L^2(\mathbb{R}^n)) . \tag{4.9}$$

Proof. Given $z_\circ \in \mathbb{C}\backslash\mathbb{R}$, $\mathrm{Re}(z_\circ) \ge \lambda_\circ$, let

$$B := (\tau(-A + \bar{z}_\circ)^{-1})' \Pi' (\Theta + \Pi M_{z_\circ} \Pi')^{-1}$$

and

$$C := \Pi \tau (-A + z_\circ)^{-1} ,$$

so that, by (4.7),

$$(-A_{\Pi,\Theta} + z_\circ)^{-1} - (-A + z_\circ)^{-1} = BC .$$

By (3.18), one has $(-A + z_\circ)^{-(m+1)} \in \mathsf{B}(L^2(\mathbb{R}^n), H^{2m+2}(\mathbb{R}^n))$ and so $\tau(-A + z_\circ)^{-(m+1)}$ has range in $H^{2m+\frac{3}{2}}(\Gamma) \oplus H^{2m+\frac{1}{2}}(\Gamma)$. Then, by [11, Lemma 4.7],

$$(-A + z_\circ)^{-(m+1)} \in \mathfrak{S}_{\frac{n-1}{2m},\infty}(L^2(\mathbb{R}^n), H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma))$$

and so

$$C(-A + z_\circ)^{-m} \in \mathfrak{S}_{\frac{n-1}{2m},\infty}(L^2(\mathbb{R}^n), \mathrm{ran}(\Pi)) .$$

By duality, that also gives

$$(-A + z_\circ)^{-m}(\tau(-A + \bar{z}_\circ)^{-1})'\Pi = (\Pi\tau(-A + \bar{z}_\circ)^{-(m+1)})' \in \mathfrak{S}_{\frac{n-1}{2m},\infty}(\mathrm{ran}(\Pi'), L^2(\mathbb{R}^n)) . \tag{4.10}$$

Since $(\Theta + \Pi M_{z_\circ} \Pi')^{-1} \in \mathsf{B}(\mathrm{ran}(\Pi), \mathrm{ran}(\Pi'))$ and its range is contained in $H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma)$, by [11, Lemma 4.7], one gets

$$(\Theta + \Pi M_{z_\circ} \Pi')^{-1} \in \mathfrak{S}_{\frac{n-1}{s},\infty}(\mathrm{ran}(\Pi), H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma)) ,$$

and so, by (4.10),

$$(-A + z_\circ)^{-m} B \in \mathfrak{S}_{\frac{n-1}{2m+s},\infty}(\mathrm{ran}(\Pi), L^2(\mathbb{R}^n)) .$$

The proof is then concluded by [12, Lemma 2.3]. □

Next we provide a version of Theorem 4.11 which applies to the case where the self-adjoint operator Θ is defined through the associated sesquilinear form. In particular the next results apply to all the operators considered in Section 6.

Theorem 4.12. *Suppose that* Γ *is smooth. Let* $\tilde f$ *be the sesquilinear form associated to the self-adjoint operator in* $\mathrm{ran}(\Pi)$ *defined by* $\tilde\Theta := \Theta(\Lambda^3 \oplus \Lambda)$*; if* $\mathrm{dom}(\tilde f) \subseteq H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma)$ *with*

$$s_* = \min\left\{s_1 - \frac{3}{2}, s_2 - \frac{1}{2}\right\} \geq 1\,,$$

then for any integer $k \geq 1$ *and for any* $z \in \rho(A) \cap \rho(A_{\Pi,\Theta})$ *one has*

$$(-A_{\Pi,\Theta} + z)^{-k} - (-A + z)^{-k} \in \mathfrak{S}_{\frac{n-1}{2(k-1)+2s_*},\infty}(L^2(\mathbb{R}^n))\,. \tag{4.11}$$

Proof. According to our assumptions,

$$\mathrm{dom}(\Theta) = (\Lambda^3 \oplus \Lambda)\mathrm{dom}(\tilde\Theta) \subseteq (\Lambda^3 \oplus \Lambda)\mathrm{dom}(\tilde f) \subseteq H^{-\frac{1}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)\,.$$

Therefore hypothesis (4.8) holds and so, by Lemma 4.10, $\sigma_{ess}(A_{\Pi,\Theta}) = \sigma_{ess}(A) \subseteq (-\infty, \|V_{\text{neg}}\|_\infty]$. Thus there exists $\lambda \in \rho(A) \cap \rho(A_{\Pi,\Theta}) \cap \mathbb{R}$ and so, by (4.7), the operator $\tilde\Theta + \Pi\tilde M_\lambda\Pi$ is self-adjoint and has a bounded inverse $(\tilde\Theta + \Pi\tilde M_\lambda\Pi)^{-1} \in \mathsf{B}(\mathrm{ran}(\Pi))$ (here $\tilde M_z := M_z(\Lambda^3 \oplus \Lambda)$). Let

$$\tilde\Theta + \Pi\tilde M_\lambda\Pi = \tilde U\,|\tilde\Theta + \Pi\tilde M_\lambda\Pi|$$

be the polar decomposition of $\tilde\Theta + \Pi\tilde M_\lambda\Pi$ (see e.g. [49, Section 7, Chapter VI]). Since $\tilde\Theta + \Pi\tilde M_\lambda\Pi$ is self-adjoint and injective, $\tilde U$ is self-adjoint and unitary. Then

$$(\tilde\Theta + \Pi\tilde M_\lambda\Pi)^{-1} = |\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-1}\tilde U = |\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}}|\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}}\tilde U\,.$$

Since $|\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}}$ is bounded and commutes with $\tilde\Theta + \Pi\tilde M_\lambda\Pi$, by [49, Lemma 2.37, Chapter VI], it commutes with $\tilde U$. Therefore

$$(\tilde\Theta + \Pi\tilde M_\lambda\Pi)^{-1} = |\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}}\tilde U\,|\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}}$$

and so, by (4.7),

$$(-A_{\Pi,\Theta} + \lambda)^{-1} - (-A + \lambda)^{-1} = BC\,,$$

where

$$B = (\tau(-A + \lambda)^{-1})^*\Pi\,|\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}}\,, \quad C = \tilde U B^*\,.$$

Since $\Pi\tilde M_\lambda\Pi$ is bounded, $\mathrm{dom}(\tilde f) = \mathrm{dom}(|\tilde\Theta|^{\frac{1}{2}}) = \mathrm{dom}(|\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{\frac{1}{2}})$ and so

$$\mathrm{ran}(|\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}}) \subseteq H^{s_1}(\Gamma) \oplus H^{s_2}(\Gamma)\,.$$

Therefore, by [11, Lemma 4.7], one gets

$$|\tilde\Theta + \Pi\tilde M_\lambda\Pi|^{-\frac{1}{2}} \in \mathfrak{S}_{\frac{n-1}{s_*},\infty}(\mathrm{ran}(\Pi), H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma))\,.$$

Then, by (4.10),

$$(-A + \lambda)^{-m}B \in \mathfrak{S}_{\frac{n-1}{2m+s_*},\infty}(\mathrm{ran}(\Pi), L^2(\mathbb{R}^n))\,.$$

and

$$C(-A + \lambda)^{-m} \in \mathfrak{S}_{\frac{n-1}{2m+s_*},\infty}(L^2(\mathbb{R}^n), \mathrm{ran}(\Pi))\,.$$

The proof is then concluded by [12, Lemma 2.3]. ☐

Corollary 4.13. *Let* Γ *be smooth and let* Θ *satisfy the same hypotheses as either in Theorem 4.11 or in Theorem 4.12. Then*

$$\sigma_{ac}(A_{\Pi,\Theta}) = \sigma_{ac}(A)$$

and the wave operators

$$W_\pm(A_{\Pi,\Theta}, A) = \text{s-}\lim_{t\to\pm\infty} e^{-itA_{\Pi,\Theta}}e^{itA}P_{ac}(A)\,,$$

$$W_\pm(A, A_{\Pi,\Theta}) = \text{s-}\lim_{t\to\pm\infty} e^{-itA}e^{itA_{\Pi,\Theta}}P_{ac}(A_{\Pi,\Theta})$$

exist and are complete, i.e. the limits exist everywhere and the ranges coincide with with the absolutely continuous subspaces.

Proof. By either Theorem 4.11 or Theorem 4.12, for k large enough, the resolvent difference $(-A_{\Pi,\Theta}+\lambda)^{-k}-(-A+\lambda)^{-k}$ is trace class; so, by the Birman-Kato criterion (see [49, Theorem 4.8, Chapter X]), one obtains the existence and completeness of the wave operators; thus $\sigma_{ac}(A_{\Pi,\Theta})=\sigma_{ac}(A)$. ☐

Remark 4.14. In the case $A\le 0$ and $A_{\Pi,\Theta}\le 0$, by Corollary 4.13 and [48, Sections 8 and 9], one also gets the existence and completeness of wave operators for the pairs of wave equations $\partial^2_{tt}u=A_{\Pi,\Theta}u$ and $\partial^2_{tt}u=Au$.

Remark 4.15. Under additional hypotheses on the behavior at infinity of the coefficients of A, the spectral results in Lemma 4.10 and Corollary 4.13 can be specified. Let us suppose that

$$a_{ij}(x)=a^\circ_{ij}+b_{ij}(x)\,,\quad b_{ij}(x)=O(1/\|x\|^\delta)\,,\tag{4.12}$$

$$\partial_{x_i}b_{ij}(x)=O(1/\|x\|^\delta)\,,\quad V(x)=O(1/\|x\|^\delta)\,,\tag{4.13}$$

for some $\delta>1$, as $\|x\|\to+\infty$. Let $A_\circ$ be the differential operator with constant coefficients

$$A_\circ: H^2(\mathbb{R}^n)\subset L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)\,,\quad A_\circ u=\sum_{1\le i,j\le n}a^\circ_{ij}\partial^2_{x_ix_j}u\,.$$

One has $\sigma(A_\circ)=\sigma_{ac}(A_\circ)=\sigma_{ess}(A_\circ)=(-\infty,0]$. Since, by [7, Theorem 5.3], $\sigma_{ess}(A)=\sigma_{ess}(A_\circ)$ and, by [44, Theorem 2.1], [45, Chapter XIV], $\sigma_{ac}(A)=\sigma_{ac}(A_\circ)$, Lemma 4.10 and Corollary 4.13 give

$$\sigma_{ess}(A_{\Pi,\Theta})=\sigma_{ac}(A_{\Pi,\Theta})=(-\infty,0]\,.$$

5. Applications: boundary conditions on Γ.

Using the scheme provided by Theorem 4.4 and Corollary 4.9, we next give the construction of some standard models of elliptic operators with boundary conditions on Γ, the boundary of a bounded domain Ω of class $C^{1,1}$.

5.1. Dirichlet boundary conditions.

Let us consider the self-adjoint extension A_D corresponding to Dirichlet boundary conditions on the whole Γ; it is given by the direct sum $A_D=A^D_-\oplus A^D_+$, where the self-adjoint operators $A^D_\pm$ are defined by $A^D_\pm:=A|\text{dom}(A^D_\pm)$, $\text{dom}(A^D_\pm)=\{u_\pm\in H^2(\Omega_\pm):\gamma^\pm_0u_\pm=0\}$. Since

$$\begin{aligned}\text{dom}(A^D_-)\oplus\text{dom}(A^D_+)=&\{u=u_-\oplus u_+\in H^2(\Omega_-)\oplus H^2(\Omega_+):[\gamma_0]u=0\,,\gamma_0u=0\}\\=&\{u\in H^1(\mathbb{R}^n)\cap H^2(\mathbb{R}^n\backslash\Gamma):\gamma_0u=0\}\,,\end{aligned}$$

that corresponds, in Corollary 4.9, to the choice $\Pi=\Pi_1$, where $\Pi_1(\phi\oplus\varphi):=\phi\oplus 0$, and $B_\Theta=0$, i.e. $\Theta(\phi\oplus\varphi):=(-\Theta_D\phi)\oplus 0$, where Θ_D is the (necessarily self-adjoint, by Lemma 2.6) operator

$$\Theta_D=\gamma_0\text{SL}:H^{\frac12}(\Gamma)\subseteq H^{-\frac32}(\Gamma)\to H^{\frac32}(\Gamma)\,.\tag{5.1}$$

Thus

$$(A^D_-\oplus A^D_+)u=Au-[\gamma_1]u\,\delta_\Gamma$$

and, for any $z\in\rho(A)\cap\rho(A^D_-)\cap\rho(A^D_+)$,

$$(-(A^D_-\oplus A^D_+)+z)^{-1}=(-A+z)^{-1}-\text{SL}_z(\gamma_0\text{SL}_z)^{-1}\gamma_0(-A+z)^{-1}\,.\tag{5.2}$$

Let $z\in\rho(A)\cap\rho(A^D_-)\cap\rho(A^D_+)$, so that, by (5.2), $(\gamma_0\text{SL}_z)^{-1}\in\mathsf{B}(H^{\frac32}(\Gamma),H^{\frac12}(\Gamma))$. Given $\varphi\in H^{\frac32}(\Gamma)$, let us define $\phi:=(\gamma_0\text{SL}_z)^{-1}\varphi$ and $u_\pm:=\text{SL}^\pm_z\phi$. Then $A^{\max}_\pm u_\pm=z\,u_\pm$ and $\gamma^\pm_0u_\pm=\varphi$ and so one gets $u_\pm=K^\pm_z\varphi$, where $K^\pm_z\in\mathsf{B}(H^s(\Gamma),\text{dom}(A^{\max}_\pm))$, $s\ge-\frac12$, is the Poisson operator which solves the Dirichlet boundary value problem

$$\begin{cases}(A^{\max}_\pm-z)K^\pm_z\psi=0\\ \hat\gamma^\pm_0K^\pm_z\psi=\psi\,.\end{cases}\tag{5.3}$$

To $K^\pm_z$ one associates the Dirichlet-to-Neumann operator $P^\pm_z\in\mathsf{B}(H^s(\Gamma),H^{s-1}(\Gamma))$, $s\ge-\frac12$, defined by $P^\pm_z:=\hat\gamma^\pm_1K^\pm_z$. Thus, since $[\gamma_1]\text{SL}_z\phi=-\phi$, one has

$$\forall z\in\rho(A)\cap\rho(A^D_-)\cap\rho(A^D_+)\,,\qquad(\gamma_0\text{SL}_z)^{-1}=P^-_z-P^+_z\,.\tag{5.4}$$

Therefore, by (5.2),

$$(-(A^D_-\oplus A^D_+)+z)^{-1}=(-A+z)^{-1}-\text{SL}_z(P^-_z-P^+_z)\gamma_0(-A+z)^{-1}\,.$$

5.2. Neumann boundary conditions.

Let us consider the self-adjoint extension A_N corresponding to Neumann boundary conditions on the whole Γ; it is given by the direct sum $A_N = A^N_- \oplus A^N_+$, where the self-adjoint operators $A^N_\pm$ are defined by $A^N_\pm := A|\mathrm{dom}(A^N_\pm)$, $\mathrm{dom}(A^N_\pm) = \{u_\pm \in H^2(\Omega_\pm) : \gamma_1^\pm u_\pm = 0\}$. Since

$$\mathrm{dom}(A^N_-) \oplus \mathrm{dom}(A^N_+) = \{u \in H^2(\mathbb{R}^n\backslash\Gamma) : [\gamma_1]u = 0\,, \gamma_1 u = 0\}\,,$$

that corresponds, in Corollary 4.9, to the choice $\Pi = \Pi_2$, where $\Pi_2(\phi\oplus\varphi) := 0\oplus\varphi$, and $B_\Theta = 0$, i.e. $\Theta(\phi\oplus\varphi) := 0\oplus(-\Theta_N\phi)$, where Θ_N is the (necessarily self-adjoint, by Lemma 2.6) operator

$$\Theta_N = \gamma_1\mathrm{DL} : H^{\frac{3}{2}}(\Gamma) \subseteq H^{-\frac{1}{2}}(\Gamma) \to H^{\frac{1}{2}}(\Gamma)\,. \tag{5.5}$$

Thus

$$(A^N_- \oplus A^N_+)u = Au - [\gamma_0]u\,\partial_{\underline{a}}\delta_\Gamma\,,$$

and, for any $z \in \rho(A) \cap \rho(A^N_-) \cap \rho(A^N_+)$

$$(-(A^N_- \oplus A^N_+) + z)^{-1} =(-A+z)^{-1} - \mathrm{DL}_z(\gamma_1\mathrm{DL}_z)^{-1}\gamma_1(-A+z)^{-1}\,. \tag{5.6}$$

Let $z \in \rho(A)\cap\rho(A^N_-)\cap\rho(A^N_+)$, so that, by (5.6), $(\gamma_1\mathrm{DL}_z)^{-1} \in \mathsf{B}(H^{\frac{1}{2}}(\Gamma), H^{-\frac{1}{2}}(\Gamma))$. Given $\phi \in H^{\frac{1}{2}}(\Gamma)$, let us define $\varphi := (\gamma_1\mathrm{DL}_z)^{-1}\phi$ and $u_\pm := \mathrm{SL}^\pm_z\varphi$. Then $A^{\max}_\pm u_\pm = z\,u_\pm$ and $\gamma^\pm_1 u_\pm = \phi$ and so one gets $u_\pm = \tilde{K}^\pm_z\phi$, where $\tilde{K}^\pm_z \in \mathsf{B}(H^s(\Gamma), \mathrm{dom}(A^{\max}_\pm))$, $s \geq -\frac{3}{2}$, solves the boundary value problem

$$\begin{cases}(A^{\max}_\pm - z)\tilde{K}^\pm_z\psi = 0\\ \hat{\gamma}^\pm_1\tilde{K}^\pm_z\psi = \psi\,.\end{cases}$$

To $\tilde{K}^\pm_z$ one associates the Neumann-to-Dirichlet operator $Q^\pm_z \in \mathsf{B}(H^s(\Gamma), H^{s+1}(\Gamma))$, $s \geq -\frac{3}{2}$, defined by $Q^\pm_z := \hat{\gamma}^\pm_0\tilde{K}^\pm_z$. Thus, since $[\gamma_0]\mathrm{DL}_z\varphi = \varphi$, one has

$$\forall z \in \rho(A) \cap \rho(A^N_-) \cap \rho(A^N_+)\,, \qquad (\gamma_1\mathrm{DL}_z)^{-1} = Q^+_z - Q^-_z\,. \tag{5.7}$$

Therefore, by (5.6),

$$(-(A^N_- \oplus A^N_+) + z)^{-1} = (-A+z)^{-1} + \mathrm{DL}_z(Q^-_z - Q^+_z)\gamma_1(-A+z)^{-1}\,.$$

5.3. Robin boundary conditions.

Let us consider the linear operator A_R corresponding to Robin boundary conditions on the whole Γ; it is given by the direct sum $A_R = A^R_- \oplus A^R_+$, where

$$A^R_\pm := A|\mathrm{dom}(A^R_\pm)\,, \qquad \mathrm{dom}(A^R_\pm) = \{u_\pm \in \mathrm{dom}(A^{\max}_\pm) : \gamma^\pm_1 u_\pm = b_\pm\,\gamma^\pm_0 u_\pm\}\,.$$

We suppose that $b_\pm \in M^{\frac{1}{2}}(\Gamma)$ and that the functions $b_\pm$ are real-valued. Hence the operators $A^R_\pm$ are self-adjoint and $\mathrm{dom}(A^\pm_R) \subseteq H^2(\Omega_\pm)$ (use e.g [41, Theorem 11]). In case $b_+(x) \neq b_-(x)$ for a.e. $x \in \Gamma$, the domain of $A^R_- \oplus A^R_+$ represents as

$$\begin{aligned}&\mathrm{dom}(A^R_- \oplus A^R_+) = \{u \in H^2(\mathbb{R}^n\backslash\Gamma) : \gamma^\pm_1 u_\pm = b_\pm\,\gamma^\pm_0 u_\pm\}\\ =&\left\{u \in H^2(\mathbb{R}^n\backslash\Gamma) : (b_+ - b_-)\gamma_0 u = [\gamma_1]u - \frac{1}{2}(b_+ + b_-)[\gamma_0]u\,,\ (b_+ - b_-)\gamma_1 u = \frac{1}{2}(b_+ + b_-)[\gamma_1]u - b_+b_-[\gamma_0]u\right\}\,.\end{aligned} \tag{5.8}$$

Then, according to Corollary 4.9, the self-adjoint operator $A^R_- \oplus A^R_+$ corresponds to the choice $\Pi = 1$ and $B_\Theta = B_R$, where

$$B_R = -\frac{1}{[b]}\begin{bmatrix}1 & \langle b\rangle\\ \langle b\rangle & b_+b_-\end{bmatrix}\,, \quad \langle b\rangle := \frac{1}{2}(b_+ + b_-)\,, \qquad [b] := b_+ - b_-\,, \tag{5.9}$$

provided that the operator

$$\Theta = -\Theta_R\,, \quad \Theta_R := \begin{bmatrix}1/[b] + \gamma_0\mathrm{SL} & \langle b\rangle/[b] + \gamma_0\mathrm{DL}\\ \langle b\rangle/[b] + \gamma_1\mathrm{SL} & b_+b_-/[b] + \gamma_1\mathrm{DL}\end{bmatrix} \tag{5.10}$$

is self-adjoint. This follows by the next lemma:

Lemma 5.1. *If* $b_\pm \in M^{\frac{3}{2}}(\Gamma)$ *are real valued and* $1/[b] \in L^\infty(\Gamma)$ *, then*

$$\Theta_R : H^{\frac{3}{2}}(\Gamma) \times H^{\frac{3}{2}}(\Gamma) \subset H^{-\frac{3}{2}}(\Gamma) \oplus H^{-\frac{1}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)$$

is self-adjoint.

Proof. Θ_R is a well-defined linear operator by (3.26) and (3.27). By (5.10) and (5.8), setting $u_\circ := u + \mathrm{SL}[\gamma_1]u - \mathrm{DL}[\gamma_0]u$, one has

$$\begin{aligned}
&\{u \in H^2(\mathbb{R}^n\backslash\Gamma) : (-[\gamma_1]u) \oplus [\gamma_0]u \in \mathrm{dom}(\Theta_R)\,,\ \gamma_0 u_\circ \oplus \gamma_1 u_\circ = -\Theta_R((-[\gamma_1]u) \oplus [\gamma_0]u)\} \\
=&\left\{u \in H^2(\mathbb{R}^n\backslash\Gamma) : \gamma_0 u = ([\gamma_1]u - \langle b\rangle[\gamma_0]u)/[b]\,,\ \gamma_1 u = (\langle b\rangle[\gamma_1]u - b_+b_-[\gamma_0]u)/[b]\right\} \\
=&\mathrm{dom}(A^R_- \oplus A^R_+)\,.
\end{aligned}$$

Since $A^R_- \oplus A^R_+$ is self-adjoint, Θ_R is self-adjoint by Lemma 2.6. ☐

Remark 5.2. By Green's formula and Ehrling's lemma (here we procede as in the proof of [12, Proposition 3.15]) one has, for $\epsilon > 0$ sufficiently small and $\frac{1}{2} < s < 1$,

$$\begin{aligned}
\langle(-A^R_\pm + \lambda_\circ)u_\pm, u_\pm\rangle_{L^2(\Omega_\pm)} \geq& c_\circ\|\nabla u_\pm\|^2_{L^2(\Omega_\pm)} - \|b_\pm\|_{L^\infty(\Gamma)}\|\gamma_0^\pm u_\pm\|^2_{L^2(\Gamma)} \\
\geq& c_\circ\|\nabla u_\pm\|^2_{L^2(\Omega_\pm)} - \|b_\pm\|_{L^\infty(\Gamma)}\|u_\pm\|^2_{H^s(\Omega_\pm)} \\
\geq& c_\circ\|\nabla u_\pm\|^2_{L^2(\Omega_\pm)} - \epsilon\,\|b_\pm\|_{L^\infty(\Gamma)}\|u_\pm\|^2_{H^1(\Omega_\pm)} - c^\pm_\epsilon\|b_\pm\|_{L^\infty(\Gamma)}\|u_\pm\|^2_{L^2(\Omega_\pm)} \\
\geq& -\kappa^\pm_\epsilon\|u_\pm\|^2_{L^2(\Omega_\pm)}\,.
\end{aligned}$$

Thus $A^R_- \oplus A^R_+$ is semibounded.

According to Lemma 5.1, the Corollary 4.9 applies and we get

$$(A^R_- \oplus A^R_+)u = Au - \frac{4}{[b]}\left((\langle b\rangle\,\gamma_1 u - b_+b_-\gamma_0 u)\,\delta_\Gamma + (\gamma_1 u - \langle b\rangle\,\gamma_0 u)\,\partial_{\underline{a}}\delta_\Gamma\right)\,.$$

Moroever, for any $z \in \rho(A) \cap \rho(A^R_-) \cap \rho(A^R_+)$,

$$(-(A^R_- \oplus A^R_+) + z)^{-1}u = (-A+z)^{-1}u - G_z\begin{bmatrix} 1/[b] + \gamma_0\mathrm{SL}_z & \langle b\rangle/[b] + \gamma_0\mathrm{DL}_z \\ \langle b\rangle/[b] + \gamma_1\mathrm{SL}_z & b_+b_-/[b] + \gamma_1\mathrm{DL}_z \end{bmatrix}^{-1}\begin{bmatrix}\gamma_0(-A+z)^{-1}u \\ \gamma_1(-A+z)^{-1}u\end{bmatrix},$$

where G_z is defined in (4.2). Let us notice that the case in which one has the same Robin boundary conditions on both sides of Γ corresponds to the choice $b_+ = b = -b_-$. Thus in this case one has

$$(A^R_- \oplus A^R_+)u = Au - 2b\,\gamma_0 u\,\delta_\Gamma - (2/b)\,\gamma_1 u\,\partial_{\underline{a}}\delta_\Gamma$$

and

$$(-(A^R_- \oplus A^R_+) + z)^{-1}u = (-A+z)^{-1}u - G_z\begin{bmatrix} 1/(2b) + \gamma_0\mathrm{SL}_z & \gamma_0\mathrm{DL}_z \\ \gamma_1\mathrm{SL}_z & -b/2 + \gamma_1\mathrm{DL}_z \end{bmatrix}^{-1}\begin{bmatrix}\gamma_0(-A+z)^{-1}u \\ \gamma_1(-A+z)^{-1}u\end{bmatrix}.$$

5.4. *δ-interactions.*

Let $\Pi(\phi\oplus\varphi) = \Pi_1(\phi\oplus\varphi) := \phi\oplus 0$ and $\Theta(\phi\oplus\varphi) = (-\Theta_{\alpha,D})\phi\oplus 0$, where $\Theta_{\alpha,D} := 1/\alpha + \Theta_D = 1/\alpha + \gamma_0\mathrm{SL}$ is the compression to $\mathrm{ran}(\Pi_1)$ of Θ_R (here we consider the case $b_+ = -b_- = \alpha/2$). This gives the boundary condition $\alpha\gamma_0 u = [\gamma_1]u$ and so one obtains the self-adjoint extensions usually called "δ-interactions on Γ" (see [12], [15] and references therein). In order to apply Corollary 4.9 we need the following

Lemma 5.3. *If* $\alpha \in M^{\frac{3}{2}}(\Gamma)$ *is real valued and* $1/\alpha \in L^\infty(\Gamma)$, *then* $\Theta_{\alpha,D} : H^{\frac{3}{2}}(\Gamma) \subseteq H^{-\frac{3}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma)$ *is self-adjoint.*

Proof. By $1/\alpha \in M^{\frac{3}{2}}(\Gamma)$, the linear operator $(1/\alpha) : H^{\frac{3}{2}}(\Gamma) \subset H^{-\frac{3}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma)$ is well-defined; by $\alpha \in M^{\frac{3}{2}}(\Gamma)$ it is not difficult to check that it is self-adjoint. By (3.26), $\gamma_0\mathrm{SL}\alpha \in \mathsf{B}(H^{\frac{1}{2}}(\Gamma), H^{\frac{3}{2}}(\Gamma))$. Thus, by (3.2), for any $\epsilon > 0$,

$$\begin{aligned}
\|\gamma_0\mathrm{SL}\phi\|_{H^{\frac{3}{2}}(\Gamma)} =&\|\gamma_0\mathrm{SL}\alpha(1/\alpha)\phi\|_{H^{\frac{3}{2}}(\Gamma)} \le c\,\|(1/\alpha)\phi\|_{H^{\frac{1}{2}}(\Gamma)}\\
\le&c\,\epsilon\,\|(1/\alpha)\phi\|_{H^{\frac{3}{2}}(\Gamma)} + c_\epsilon\,\|(1/\alpha)\phi\|_{H^{-\frac{3}{2}}(\Gamma)}\\
\le&c\left(\epsilon\,\|(1/\alpha)\phi\|_{H^{\frac{3}{2}}(\Gamma)} + c_\epsilon\,\|\phi\|_{H^{-\frac{3}{2}}(\Gamma)}\right)
\end{aligned}$$

and so the self-adjoint operator $\gamma_0\mathrm{SL} : H^{\frac{1}{2}}(\Gamma) \subset H^{-\frac{3}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma)$ is infinitesimally $(1/\alpha)$-bounded. The proof is then concluded by [43, Corollary 1]. ☐

Therefore, by Corollary 4.9 one gets the self-adjoint extension

$$A_{\alpha,\delta}\,u = Au - \alpha\gamma_0 u\,\delta_\Gamma\,,$$

$$\mathrm{dom}(A_{\alpha,\delta}) := \{u \in H^1(\mathbb{R}^n) \cap H^2(\mathbb{R}^n\backslash\Gamma) : \alpha\gamma_0 u = [\gamma_1]u\}\,.$$

By (4.9) and (5.4), its resolvent is given by

$$\begin{aligned}
(-A_{\alpha,\delta} + z)^{-1} =& (-A + z)^{-1} - \mathrm{SL}_z((1/\alpha) + \gamma_0\mathrm{SL}_z)^{-1}\gamma_0(-A + z)^{-1}\\
=&(-A + z)^{-1} - \mathrm{SL}_z(P_z^- - P_z^+)(\alpha + P_z^- - P_z^+)^{-1}\alpha\gamma_0(-A + z)^{-1}\,.
\end{aligned}$$

5.5. δ'-interactions.

Let $\Pi(\phi \oplus \varphi) = \Pi_2(\phi \oplus \varphi) := 0 \oplus \varphi$ and $\Theta(\phi \oplus \varphi) = 0 \oplus (-\Theta_{\beta,N}\varphi)$, where $\Theta_{\beta,N} := -1/\beta + \Theta_N = -1/\beta + \gamma_1\mathrm{DL}$ is the compression to $\mathrm{ran}(\Pi_2)$ of Θ_R (here we consider the case $b_+ = -b_- = 2/\beta$). This gives the boundary condition $\beta\gamma_1 u = [\gamma_0]u$ and so one obtains the self-adjoint extensions usually called "δ'-interactions on Γ" (see [12]and references therein). In order to apply Corollary 4.9 we need the following

Lemma 5.4. *If $\beta \in M^{\frac{1}{2}}(\Gamma)$ is real valued and $1/\beta \in L^\infty(\Gamma)$, then $\Theta_{\beta,N} : H^{\frac{3}{2}}(\Gamma) \subseteq H^{-\frac{1}{2}}(\Gamma) \to H^{\frac{1}{2}}(\Gamma)$ is self-adjoint.*

Proof. By (3.30), one has $(\hat\gamma_1\mathrm{DL})^{-1} \in \mathsf{B}(H^{-\frac{1}{2}}(\Gamma), H^{\frac{1}{2}}(\Gamma))$. Thus, by (3.2) and (3.27), one gets, for any $\epsilon > 0$,

$$\begin{aligned}
\|(1/\beta)\varphi\|_{H^{\frac{1}{2}}(\Gamma)} =&\|(1/\beta)(\hat\gamma_1\mathrm{DL})^{-1}\hat\gamma_1\mathrm{DL}\varphi\|_{H^{\frac{1}{2}}(\Gamma)} \le c\,\|\hat\gamma_1\mathrm{DL}\varphi\|_{H^{-\frac{1}{2}}(\Gamma)}\\
\le&c\,\epsilon\,\|\hat\gamma_1\mathrm{DL}\varphi\|_{H^{\frac{1}{2}}(\Gamma)} + c_\epsilon\,\|\hat\gamma_1\mathrm{DL}\varphi\|_{H^{-\frac{3}{2}}(\Gamma)}\\
\le&c\left(\epsilon\,\|\hat\gamma_1\mathrm{DL}\varphi\|_{H^{\frac{1}{2}}(\Gamma)} + c_\epsilon\,\|\varphi\|_{H^{-\frac{1}{2}}(\Gamma)}\right)
\end{aligned}$$

and so the operator $(1/\beta) : H^{\frac{3}{2}}(\Gamma) \subseteq H^{-\frac{1}{2}}(\Gamma) \to H^{\frac{1}{2}}(\Gamma)$ is infinitesimally $\hat\gamma_1\mathrm{DL}$-bounded. Since $\hat\gamma_1\mathrm{DL} : H^{\frac{3}{2}}(\Gamma) \subseteq H^{-\frac{1}{2}}(\Gamma) \to H^{\frac{1}{2}}(\Gamma)$ is self-adjoint, the proof is then concluded by [43, Corollary 1]. ☐

Therefore, by Corollary 4.9 one gets the self-adjoint extension

$$A_{\beta,\delta'}u = Au - \beta\gamma_1 u\,\partial_{\underline{a}}\delta_\Gamma\,,$$

$$\mathrm{dom}(A_{\beta,\delta'}) := \{u \in H^2(\mathbb{R}^n\backslash\Gamma) : [\gamma_1]u = 0\,,\ \beta\gamma_1 u = [\gamma_0]u\}\,.$$

By (4.9) and (5.7), its resolvent is given by

$$\begin{aligned}
(-A_{\beta,\delta'} + z)^{-1} =& (-A + z)^{-1} + \mathrm{DL}_z((1/\beta) - \gamma_1\mathrm{DL}_z)^{-1}\gamma_1(-A + z)^{-1}\\
=&(-A + z)^{-1} + \mathrm{DL}_z(Q_z^+ - Q_z^-)(-\beta + Q_z^+ - Q_z^-)^{-1}\beta\gamma_1(-A + z)^{-1}\,.
\end{aligned}$$

6. Applications: boundary conditions on $\Sigma \subset \Gamma$.

Using the scheme provided by Theorem 4.4, we next give the construction of some models of elliptic operators with boundary conditions on a relatively open part $\Sigma \subset \Gamma$ of class $C^{0,1}$. In such cases we need more regularity hypotheses on Ω (with respect to the ones used in the previous section); these are needed in the proofs, in order that Sobolev spaces of appropriate order can be properly defined.

6.1. Dirichlet boundary conditions.

Given $\Sigma \subset \Gamma$ relatively open of class $C^{0,1}$, we denote by Π_Σ the orthogonal projector in the Hilbert space $H^{\frac{3}{2}}(\Gamma)$ such that $\text{ran}(\Pi_\Sigma) = H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp$. By (3.4), $\text{ran}(\Pi'_\Sigma) = H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)$, where $\Pi'_\Sigma = \Lambda^3\Pi_\Sigma\Lambda^{-3}$ is the dual projection. In the following, we also use the identifications $H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp \simeq H^{\frac{3}{2}}(\Sigma)$ and $H^{\frac{3}{2}}(\Sigma)' \simeq H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)$. We denote the orthogonal projection from $H^{\frac{3}{2}}(\Gamma)$ onto $H^{\frac{3}{2}}(\Sigma)$ by R_Σ; it can be identified with the restriction map $R_\Sigma : H^s(\Gamma) \to H^s(\Sigma)$, $R_\Sigma\phi := \phi|\Sigma$ (see the end of Subsection 3.1).

Theorem 6.1. *Let Ω be of class $C^{3,1}$, then*

$$\Theta_{D,\Sigma} : \text{dom}(\Theta_{D,\Sigma}) \subseteq H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{\frac{3}{2}}(\Sigma)\,, \quad \Theta_{D,\Sigma}\phi := R_\Sigma\gamma_0\text{SL}\Pi'_\Sigma\phi \equiv (\gamma_0\text{SL}\phi)|\Sigma\,,$$

$$\text{dom}(\Theta_{D,\Sigma}) := \{\phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : (\gamma_0\text{SL}\phi)|\Sigma \in H^{\frac{3}{2}}(\Sigma)\}$$

is self-adjoint.

Proof. Let f_D be the densely defined sesquilinear form in the Hilbert space $H^{\frac{3}{2}}(\Gamma)$

$$f_D : H^{5/2}(\Gamma) \times H^{5/2}(\Gamma) \subseteq H^{\frac{3}{2}}(\Gamma) \times H^{\frac{3}{2}}(\Gamma) \to \mathbb{R}$$

$$f_D(\phi_1, \phi_2) := \langle \Lambda^3\phi_1, \gamma_0\text{SL}\Lambda^3\phi_2\rangle_{-\frac{1}{2},\frac{1}{2}} = \langle \gamma_0\text{SL}\Lambda^3\phi_1, \Lambda^3\phi_2\rangle_{\frac{1}{2},-\frac{1}{2}}\,.$$

By (3.29),

$$f_D(\phi, \phi) \geq c_0\,\|\Lambda^3\phi\|^2_{-\frac{1}{2}} = c_0\,\|\phi\|^2_{H^{5/2}(\Gamma)}$$

and so f_D is strictly positive and closed. Since, for any $\phi_1 \in H^{7/2}(\Gamma)$ and for any $\phi_2 \in H^{5/2}(\Gamma)$,

$$f_D(\phi_1, \phi_2) = \langle \Lambda^{\frac{3}{2}}\gamma_0\text{SL}\Lambda^3\phi_1, \Lambda^{\frac{3}{2}}\phi_2, \rangle_{L^2(\Gamma)} = \langle \gamma_0\text{SL}\Lambda^3\phi_1, \phi_2\rangle_{H^{\frac{3}{2}}(\Gamma)}\,,$$

f_D is the sesquilinear form associated with the self-adjoint operator in $H^{\frac{3}{2}}(\Gamma)$ defined by

$$\tilde{\Theta}_D := \gamma_0\text{SL}\Lambda^3 : H^{7/2}(\Gamma) \subseteq H^{\frac{3}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma)\,.$$

The domain

$$\text{dom}(f_D) \cap \text{ran}(\Pi_\Sigma) = H^{5/2}(\Gamma) \cap H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp = H^{5/2}(\Gamma) \cap \Lambda^{-3}H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) = \Lambda^{-3}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$$

is dense in $\Lambda^{-3}H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)$, and we can use Lemma 7.1 to determine the positive self-adjoint operator $\tilde{\Theta}_{D,\Sigma}$ in $\text{ran}(\Pi_\Sigma)$ associated to the restriction of f_D to $\text{ran}(\Pi_\Sigma)$. Then

$$\Theta_{D,\Sigma} := \text{U}_\Sigma\tilde{\Theta}_{D,\Sigma}\Lambda^{-3}\,, \quad \text{dom}(\Theta_{D,\Sigma}) := \Lambda^3\text{dom}(\tilde{\Theta}_{D,\Sigma})\,,$$

is self-adjoint, where the map $\text{U}_\Sigma(\Pi_\Sigma\phi) = \phi|\Sigma$ provides the unitary isomorphism $H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp \simeq H^{\frac{3}{2}}(\Sigma)$. To conclude the proof we need to determine the operator $\breve{\Theta}_D := (\tilde{\Theta}_D)\breve{}$ and the subspace $K_\Sigma := \mathfrak{k}_{\Pi_\Sigma}$ (we refer to the Appendix for the notations). Let H_D be the Hilbert space given by $\text{dom}(f_D) = H^{5/2}(\Gamma)$ endowed with the scalar product $\langle\phi_1, \phi_2\rangle_D := f_D(\phi_1, \phi_2)$; let H'_D denote its dual space. Since

$$f_D(\phi_1, \phi_2) = \langle \gamma_0\text{SL}\Lambda^3\phi_1, \Lambda^3\phi_2\rangle_{\frac{1}{2},-\frac{1}{2}} = \langle \Lambda^{\frac{1}{2}}\gamma_0\text{SL}\Lambda^3\phi_1, \Lambda^{5/2}\phi_2\rangle_{L^2(\Gamma)}\,,$$

one has

$$H'_D = H^{\frac{1}{2}}(\Gamma)\,, \quad \langle\varphi, \phi\rangle_{H'_D,H_D} = \langle \Lambda^{\frac{1}{2}}\varphi, \Lambda^{5/2}\phi\rangle_{L^2(\Gamma)}$$

and

$$\breve{\Theta}_D : H^{5/2}(\Gamma) \to H^{\frac{1}{2}}(\Gamma)\,, \quad \breve{\Theta}_D = \gamma_0\text{SL}\Lambda^3\,.$$

Moreover

$$\begin{aligned}
K_\Sigma =&\{\varphi \in H^{\frac{1}{2}}(\Gamma) : \forall \phi \in \Lambda^{-3} H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) , \quad \langle \Lambda^{\frac{1}{2}}\varphi, \Lambda^{5/2}\phi\rangle_{L^2(\Gamma)} = 0\} \\
=&\{\varphi \in H^{\frac{1}{2}}(\Gamma) : \forall \phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) , \langle \Lambda^{-\frac{1}{2}}\phi, \Lambda^{\frac{1}{2}}\varphi\rangle_{L^2(\Gamma)} = 0\} \\
=&\{\varphi \in H^{\frac{1}{2}}(\Gamma) : \forall \phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) , \langle \phi, \varphi\rangle_{-\frac{1}{2},\frac{1}{2}} = 0\} \\
=&H^{\frac{1}{2}}_{\Sigma^c}(\Gamma) .
\end{aligned}$$

Therefore, by Lemma 7.1, $\tilde{\Theta}_{D,\Sigma}$ is self-adjoint on the domain

$$\mathrm{dom}(\tilde{\Theta}_{D,\Sigma}) := \{\phi \in \Lambda^{-3} H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists \tilde{\phi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } \gamma_0 \mathrm{SL}\Lambda^3\phi - \tilde{\phi} \in H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)\}$$

and $\tilde{\Theta}_{D,\Sigma}\phi = \tilde{\phi}$. Then $\mathrm{U}_\Sigma(\tilde{\Theta}_{D,\Sigma}\Lambda^{-3}\phi) = \tilde{\phi}|\Sigma = (\hat{\gamma}_0\mathrm{SL}\phi)|\Sigma$ and

$$\begin{aligned}
\mathrm{dom}(\Theta_{D,\Sigma}) =&\{\phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists \tilde{\phi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } \gamma_0 \mathrm{SL}\phi - \tilde{\phi} \in H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)\} \\
\subseteq&\{\phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : (\gamma_0 \mathrm{SL}\phi)|\Sigma \in H^{\frac{3}{2}}(\Sigma)\} \\
\subseteq&\{\phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists \tilde{\phi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } (\gamma_0 \mathrm{SL}\phi)|\Sigma = \tilde{\phi}|\Sigma\} \\
=&\{\phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists \tilde{\phi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } \gamma_0 \mathrm{SL}\phi - \tilde{\phi} \in H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)\} = \mathrm{dom}(\Theta_{D,\Sigma}) .
\end{aligned}$$

□

Corollary 6.2. *The linear operator in $L^2(\mathbb{R}^n)$ defined by $A_{D,\Sigma} := (A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{D,\Sigma})$, where*

$$\begin{aligned}
\mathrm{dom}(A_{D,\Sigma}) =&\{u \in H^1(\mathbb{R}^n) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\hat{\gamma}_1]u \in \mathrm{dom}(\Theta_{D,\Sigma}) ,\ (\gamma_0 u)|\Sigma = 0\} \\
\subseteq&\{u \in H^1(\mathbb{R}^n) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : (\gamma^\pm_0 u)|\Sigma = 0 ,\ ([\hat{\gamma}_1]u)|\overline{\Sigma}^c = 0\} ,
\end{aligned}$$

is self-adjoint and its resolvent is given by

$$(-A_{D,\Sigma} + z)^{-1} =(-A + z)^{-1} - \mathrm{SL}_z\Pi'_\Sigma(R_\Sigma\gamma_0\mathrm{SL}_z\Pi'_\Sigma)^{-1}R_\Sigma\gamma_0(-A + z)^{-1} ,$$

where R_Σ is the restriction operator $R_\Sigma\phi = \phi|\Sigma$ and Π'_Σ acts there as the inclusion map $\Pi'_\Sigma : H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{-\frac{3}{2}}(\Gamma)$.

Proof. By Theorem 6.1 and Theorem 4.4, taking $\Pi(\phi\oplus\varphi) = \Pi_\Sigma\phi\oplus 0$ and $\Theta(\phi\oplus\varphi) = (-\mathrm{U}^{-1}_\Sigma\Theta_{D,\Sigma}\phi)\oplus 0$ one gets the self-adjoint extension $(A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{D,\Sigma})$ with domain (contained in $H^1(\mathbb{R}^n\backslash\Gamma)$ by $\mathrm{dom}(\Theta_{D,\Sigma}) \subseteq H^{-\frac{1}{2}}(\Gamma)$ and Remark 4.7),

$$\begin{aligned}
&\mathrm{dom}(A_{D,\Sigma}) \\
=&\{u \in H^1(\mathbb{R}^n\backslash\Gamma) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u = 0 ,\ [\hat{\gamma}_1]u \in \mathrm{dom}(\Theta_{D,\Sigma}) ,\ \Pi_\Sigma\gamma_0(u + \mathrm{SL}[\hat{\gamma}_1]u) = \mathrm{U}^{-1}_\Sigma\Theta_{D,\Sigma}[\hat{\gamma}_1]u\} \\
=&\{u \in H^1(\mathbb{R}^n) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\hat{\gamma}_1]u \in \mathrm{dom}(\Theta_{D,\Sigma}) ,\ (\gamma_0 u)|\Sigma + (\gamma_0\mathrm{SL}[\hat{\gamma}_1]u)|\Sigma = (\gamma_0\mathrm{SL}[\hat{\gamma}_1]u)|\Sigma\} \\
=&\{u \in H^1(\mathbb{R}^n) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\hat{\gamma}_1]u \in \mathrm{dom}(\Theta_{D,\Sigma}) ,\ (\gamma_0 u)|\Sigma = 0\} .
\end{aligned}$$

The formula giving $(-A_{D,\Sigma} + z)^{-1}$ is consequence of (4.7), since

$$(-\mathrm{U}^{-1}_\Sigma\Theta_{D,\Sigma} + \Pi_\Sigma\gamma_0(\mathrm{SL} - \mathrm{SL}_z)\Pi'_\Sigma)^{-1}\Pi_\Sigma = (-\Theta_{D,\Sigma} + R_\Sigma\gamma_0(\mathrm{SL} - \mathrm{SL}_z)\Pi'_\Sigma)^{-1}R_\Sigma$$

and

$$-\Theta_{D,\Sigma}\phi + R_\Sigma\gamma_0(\mathrm{SL} - \mathrm{SL}_z)\phi = -\gamma_0\mathrm{SL}\phi|\Sigma + \gamma_0\mathrm{SL}\phi|\Sigma - \gamma_0\mathrm{SL}_z\phi|\Sigma = -\gamma_0\mathrm{SL}_z\phi|\Sigma ,$$

□

Remark 6.3. Since $\mathrm{supp}([\gamma_1]u) \subseteq \overline{\Sigma}$ for any $u \in \mathrm{dom}(A_{D,\Sigma})$, one has $[\hat{\gamma}_1]u\,\delta_\Gamma = [\hat{\gamma}_1]u\,\delta_{\overline{\Sigma}}$; thus

$$A_{D,\Sigma}u = Au - [\hat{\gamma}_1]u\,\delta_{\overline{\Sigma}}$$

and so $(A_{D,\Sigma}u)|\overline{\Sigma}^c = (Au)|\overline{\Sigma}^c$. This also shows that $A_{D,\Sigma}$ is a self-adjoint extension of the symmetric operator $A|C^\infty_{\mathrm{comp}}(\mathbb{R}^n\backslash\overline{\Sigma})$. Hence it depends only on Σ and not on the whole Γ: one would obtain the same operator by considering any other bounded domain $\Omega_\circ$ with boundary $\Gamma_\circ$ such that $\Sigma \subset \Gamma_\circ$.

Remark 6.4. According to the definition of $\mathrm{dom}(\Theta_{D,\Sigma})$, the Lemma 4.8 applies and one gets

$$\mathrm{dom}(A_{D,\Sigma}) \subseteq H^2(\mathbb{R}^n \backslash (\overline{\Sigma} \cup (\partial\Sigma)^{\epsilon}))\,.$$

Remark 6.5. By the results contained in the proof of Theorem 6.1, for the domain of the sesquilinear form $f_{D,\Sigma}$ associated to the self-adjoint operator $\tilde{\Theta}_{D,\Sigma}$ one has the relation $\mathrm{dom}(f_{D,\Sigma}) \subseteq H^{5/2}(\Gamma)$. Therefore Theorem 4.12 applies and so, by Corollary 4.13, $\sigma_{ac}(A_{D,\Sigma}) = \sigma_{ac}(A)$ and the wave operators $W_\pm(A_{D,\Sigma}, A)$, $W_\pm(A, A_{D,\Sigma})$ exist and are complete.

Remark 6.6. In case $n = 3$ and both Γ and Σ are smooth, by [18, Theorem 2.4], one has that

$$(\hat{\gamma}_0 \mathrm{SL}\phi)|\Sigma \in H^{s+1}(\Sigma) \iff \phi \in H^s_{\overline{\Sigma}}(\Gamma)\,, \qquad -1 < s < 0\,. \tag{6.1}$$

Therefore $\mathrm{dom}(\Theta_{D,\Sigma}) \subseteq H^s(\Gamma)$, $-\frac{1}{2} \le s < 0$, and so, by Remark 4.7 and (3.11),

$$\mathrm{dom}(A_{D,\Sigma}) \subseteq H^{\frac{3}{2}-}(\mathbb{R}^3)\,.$$

6.2. Neumann boundary conditions.

Given $\Sigma \subset \Gamma$ relatively open of class $C^{0,1}$, here we denote by Π_Σ the orthogonal projector in the Hilbert space $H^{\frac{1}{2}}(\Gamma)$ such that $\mathrm{ran}(\Pi_\Sigma) = H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp$. By (3.4), $\mathrm{ran}(\Pi'_\Sigma) = H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$, where $\Pi'_\Sigma = \Lambda\Pi_\Sigma\Lambda^{-1}$ is the dual projection. In the following, we also use the identifications $H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp \simeq H^{\frac{1}{2}}(\Sigma)$ and $H^{\frac{1}{2}}(\Sigma)' \simeq H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$. We denote the orthogonal projection from $H^{\frac{1}{2}}(\Gamma)$ onto $H^{\frac{1}{2}}(\Sigma)$ by R_Σ; it can be identified with the restriction map $R_\Sigma : H^s(\Gamma) \to H^s(\Sigma)$, $R_\Sigma\phi := \phi|\Sigma$ (see the end of Subsection 3.1).

Similarly to Example 6.1 one has the following

Theorem 6.7. *Let Ω be of class $C^{2,1}$, then*

$$\Theta_{N,\Sigma} : \mathrm{dom}(\Theta_{N,\Sigma}) \subseteq H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{\frac{1}{2}}(\Sigma)\,, \quad \Theta_{N,\Sigma} := R_\Sigma\hat{\gamma}_1 \mathrm{DL}\Pi'_\Sigma\varphi \equiv (\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma\,,$$

$$\mathrm{dom}(\Theta_{N,\Sigma}) := \{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : (\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma)\}\,,$$

is self-adjoint.

Proof. Let f_N be the densely defined sesquilinear form in the Hilbert space $H^{\frac{1}{2}}(\Gamma)$

$$f_N : H^{\frac{3}{2}}(\Gamma) \times H^{\frac{3}{2}}(\Gamma) \subseteq H^{\frac{1}{2}}(\Gamma) \times H^{\frac{1}{2}}(\Gamma) \to \mathbb{R}$$

$$f_N(\varphi_1, \varphi_2) := \langle \hat{\gamma}_1 \mathrm{DL}\Lambda\varphi_1, \Lambda\varphi_2\rangle_{-\frac{1}{2},\frac{1}{2}}\,.$$

By (3.30),

$$-f_N(\varphi, \varphi) \ge c_1\,\|\Lambda\varphi\|^2_{\frac{1}{2}} = c_1\,\|\varphi\|^2_{H^{\frac{3}{2}}(\Gamma)}$$

and so f_N is strictly negative and closed. Since, for any $\varphi_1 \in H^{5/2}(\Gamma)$ and for any $\phi_2 \in H^{\frac{3}{2}}(\Gamma)$,

$$f_N(\varphi_1, \varphi_2) = \langle \Lambda^{\frac{1}{2}}\hat{\gamma}_1\mathrm{DL}\Lambda\varphi_1, \Lambda^{\frac{1}{2}}\varphi_2, \rangle_{L^2(\Gamma)} = \langle \hat{\gamma}_1\mathrm{DL}\Lambda\varphi_1, \varphi_2\rangle_{H^{\frac{1}{2}}(\Gamma)}\,,$$

f_N is the sesquilinear form associated with the self-adjoint operator

$$\tilde{\Theta}_N := \hat{\gamma}_1\mathrm{DL}\Lambda : H^{5/2}(\Gamma) \subset H^{\frac{1}{2}}(\Gamma) \to H^{\frac{1}{2}}(\Gamma)\,.$$

Since

$$\mathrm{dom}(f_N) \cap \mathrm{ran}(\Pi_\Sigma) = H^{\frac{3}{2}}(\Gamma) \cap H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp = H^{\frac{3}{2}}(\Gamma) \cap \Lambda^{-1}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) = \Lambda^{-1}H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$$

is dense in $\Lambda^{-1}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$, we can use Lemma 7.1 to determine the positive self-adjoint operator $\tilde{\Theta}_{N,\Sigma}$ in $\mathrm{ran}(\Pi_\Sigma)$ associated to the restriction of f_N to $\mathrm{ran}(\Pi_\Sigma)$. Then

$$\Theta_{N,\Sigma} := \mathrm{U}_\Sigma\tilde{\Theta}_{N,\Sigma}\Lambda^{-1}\,, \qquad \mathrm{dom}(\Theta_{N,\Sigma}) := \Lambda\mathrm{dom}(\tilde{\Theta}_{N,\Sigma})\,,$$

is self-adjoint, where the map $\mathrm{U}_\Sigma(\Pi_\Sigma)\varphi = \varphi|\Sigma$ provides the unitary isomorphism $H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp \simeq H^{\frac{1}{2}}(\Sigma)$. To conclude the proof we need to determine the operator $\breve{\Theta}_N := (\tilde{\Theta}_N)\breve{}$ and the subspace $K_\Sigma := \mathfrak{k}_{\Pi_\Sigma}$ (we refer to the Appendix for the notations). Let H_N be the Hilbert space given by $\mathrm{dom}(f_N) = H^{\frac{3}{2}}(\Gamma)$ endowed with the scalar product $\langle\varphi_1, \varphi_2\rangle_N := -f_N(\varphi_1, \varphi_2)$; let H'_N denote its dual space. Since

$$f_N(\varphi_1, \varphi_2) = \langle\hat{\gamma}_1 \mathrm{DL}\Lambda\varphi_1, \Lambda\varphi_2\rangle_{-\frac{1}{2},\frac{1}{2}} = \langle\Lambda^{-\frac{1}{2}}\hat{\gamma}_1\mathrm{DL}\Lambda\varphi_1, \Lambda^{\frac{3}{2}}\varphi_2\rangle_{L^2(\Gamma)}\,,$$

one has

$$H'_N = H^{-\frac{1}{2}}(\Gamma)\,, \quad \langle\phi, \varphi\rangle_{H'_N,H_N} = \langle\Lambda^{-\frac{1}{2}}\phi, \Lambda^{\frac{3}{2}}\varphi\rangle_{L^2(\Gamma)}$$

and

$$\breve{\Theta}_N : H^{\frac{3}{2}}(\Gamma) \to H^{-\frac{1}{2}}(\Gamma)\,, \quad \breve{\Theta}_N = \hat{\gamma}_1\mathrm{DL}\Lambda\,.$$

Moreover

$$\begin{aligned} K_\Sigma =&\{\phi \in H^{-\frac{1}{2}}(\Gamma) : \forall\varphi \in \Lambda^{-1}H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)\,, \quad \langle\Lambda^{-\frac{1}{2}}\phi, \Lambda^{\frac{3}{2}}\varphi\rangle_{L^2(\Gamma)} = 0\}\\ =&\{\phi \in H^{-\frac{1}{2}}(\Gamma) : \forall\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)\,,\ \langle\Lambda^{-\frac{1}{2}}\phi, \Lambda^{\frac{1}{2}}\varphi\rangle_{L^2(\Gamma)} = 0\}\\ =&\{\phi \in H^{-\frac{1}{2}}(\Gamma) : \forall\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)\,,\ \langle\phi, \varphi\rangle_{-\frac{1}{2},\frac{1}{2}} = 0\}\\ =&H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\,. \end{aligned}$$

Therefore, by Lemma 5.1, $\tilde{\Theta}_{N,\Sigma}$ is self-adjoint on the domain

$$\mathrm{dom}(\tilde{\Theta}_{N,\Sigma}) := \{\varphi \in \Lambda^{-1}H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\tilde{\varphi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } \hat{\gamma}_1\mathrm{DL}\Lambda\varphi - \tilde{\varphi} \in H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\}$$

and $\tilde{\Theta}_{N,\Sigma}\varphi := \tilde{\varphi}$. Then $U_\Sigma\tilde{\Theta}_{N,\Sigma}\Lambda^{-1}\varphi = \tilde{\varphi}|\Sigma = (\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma$ and

$$\begin{aligned} \mathrm{dom}(\tilde{\Theta}_{N,\Sigma}) =&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\tilde{\varphi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } \hat{\gamma}_1\mathrm{DL}\varphi - \tilde{\varphi} \in H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\}\\ \subseteq&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : (\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma)\}\\ \subseteq&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\tilde{\varphi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } (\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma = \tilde{\varphi}|\Sigma\}\\ =&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\tilde{\varphi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } \hat{\gamma}_1\mathrm{DL}\varphi - \tilde{\varphi} \in H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\} = \mathrm{dom}(\tilde{\Theta}_{N,\Sigma}) \end{aligned}$$

□

Corollary 6.8. *The linear operator in* $L^2(\mathbb{R}^n)$ *defined by* $A_{N,\Sigma} := (A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{N,\Sigma})$, *where*

$$\begin{aligned} \mathrm{dom}(A_{N,\Sigma}) =&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{N,\Sigma})\,,\ [\hat{\gamma}_1]u = 0\,,\ (\hat{\gamma}_1 u)|\Sigma = 0\}\\ \subseteq&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : (\hat{\gamma}^\pm_1 u)|\Sigma = 0\,,\ ([\hat{\gamma}_1]u)|\overline{\Sigma}^c = 0\}\,, \end{aligned}$$

is self-adjoint and its resolvent is given by

$$(-A_{N,\Sigma} + z)^{-1} = (-A + z)^{-1} - \mathrm{DL}_z\Pi'_\Sigma(R_\Sigma\hat{\gamma}_1\mathrm{DL}_z\Pi'_\Sigma)^{-1}R_\Sigma\gamma_1(-A + z)^{-1}\,,$$

where R_Σ *is the restriction operator* $R_\Sigma\varphi = \varphi|\Sigma$ *and* Π'_Σ *acts there as the inclusion map* $\Pi'_\Sigma : H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{-\frac{1}{2}}(\Gamma)$.

Proof. By Theorem 6.1 and Theorem 4.4, taking $\Pi(\phi\oplus\varphi) = 0\oplus\Pi_\Sigma\varphi$ and $\Theta(\phi\oplus\varphi) = 0\oplus(-U^{-1}_\Sigma\Theta_{N,\Sigma}\varphi)$, one gets the self-adjoint extension $(A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{N,\Sigma})$ with domain (contained in $H^1(\mathbb{R}^n\backslash\Gamma)$ by $\mathrm{dom}(\Theta_{N,\Sigma}) \subseteq H^{\frac{1}{2}}(\Gamma)$ and Remark 4.7),

$$\begin{aligned} &\mathrm{dom}(A_{N,\Sigma})\\ =&\{u \in H^1(\mathbb{R}^n\backslash\Gamma) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{N,\Sigma})\,,\ [\hat{\gamma}_1]u = 0\,,\ \Pi_\Sigma\gamma_1(u - \mathrm{DL}[\gamma_0 u]) = -U^{-1}_\Sigma\Theta_{N,\Sigma}[\gamma_0]u\}\\ =&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{N,\Sigma})\,,\ [\hat{\gamma}_1]u = 0\,,\ (\hat{\gamma}_1 u)|\Sigma - (\hat{\gamma}_1\mathrm{DL}[\gamma_0 u])|\Sigma = -(\hat{\gamma}_1\mathrm{DL}[\gamma_0 u])|\Sigma\}\\ =&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{N,\Sigma})\,,\ [\hat{\gamma}_1]u = 0\,,\ (\hat{\gamma}_1 u)|\Sigma = 0\}\,. \end{aligned}$$

The formula giving $(-A_{N,\Sigma}+z)^{-1}$ is consequence of (4.7), since

$$(-U_{\Sigma}^{-1}\Theta_{N,\Sigma}+\Pi_{\Sigma}\gamma_1(\mathrm{DL}-\mathrm{DL}_z)\Pi_{\Sigma}')^{-1}\Pi_{\Sigma}=(-\Theta_{N,\Sigma}+R_{\Sigma}\gamma_1(\mathrm{DL}-\mathrm{DL}_z)\Pi_{\Sigma}')^{-1}R_{\Sigma}$$

and

$$-\Theta_{N,\Sigma}\varphi+R_{\Sigma}\gamma_1(\mathrm{DL}-\mathrm{DL}_z)\varphi=-(\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma+(\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma-(\hat{\gamma}_1\mathrm{DL}_z\varphi)|\Sigma=-(\hat{\gamma}_1\mathrm{DL}_z\varphi)|\Sigma\,.$$

$\square$

Remark 6.9. Since $\operatorname{supp}([\gamma_0]u)\subseteq\overline{\Sigma}$ for any $u\in\operatorname{dom}(A_{N,\Sigma})$, one has $[\gamma_0]u\,\partial_{\underline{a}}\delta_{\Gamma}=[\gamma_0]u\,\partial_{\underline{a}}\delta_{\overline{\Sigma}}$; thus

$$A_{N,\Sigma}u=Au-[\gamma_0]u\,\partial_{\underline{a}}\delta_{\overline{\Sigma}}$$

and so $(A_{N,\Sigma}u)|\overline{\Sigma}^c=(Au)|\overline{\Sigma}^c$. This also shows that $A_{N,\Sigma}$ is a self-adjoint extension of the symmetric operator $A|C^{\infty}_{\mathrm{comp}}(\mathbb{R}^n\backslash\overline{\Sigma})$. Hence it depends only on Σ and not on the whole Γ: one would obtain the same operator by considering any other bounded domain $\Omega_{\circ}$ with boundary $\Gamma_{\circ}$ such that $\Sigma\subset\Gamma_{\circ}$.

Remark 6.10. According to the definition of $\operatorname{dom}(\Theta_{N,\Sigma})$, the Lemma 4.8 applies and one gets

$$\operatorname{dom}(A_{N,\Sigma})\subseteq H^2(\mathbb{R}^n\backslash(\overline{\Sigma}\cup(\partial\Sigma)^{\epsilon}))\,.$$

Remark 6.11. By the results contained in the proof of Theorem 6.7, for the domain of the sesquilinear form $f_{N,\Sigma}$ associated to the self-adjoint operator $\tilde{\Theta}_{N,\Sigma}$ one has the relation $\operatorname{dom}(f_{N,\Sigma})\subseteq H^{\frac{3}{2}}(\Gamma)$. Therefore Theorem 4.12 applies and so, by Corollary 4.13, $\sigma_{ac}(A_{N,\Sigma})=\sigma_{ac}(A)$ and the wave operators $W_{\pm}(A_{N,\Sigma},A)$, $W_{\pm}(A,A_{N,\Sigma})$ exist and are complete.

Remark 6.12. In case $n=3$ and both Γ and Σ are smooth, by [18, Theorem 2.4], one has that

$$(\hat{\gamma}_1\mathrm{DL}\phi)|\Sigma\in H^{s-1}(\Sigma)\iff\phi\in H^{s}_{\overline{\Sigma}}(\Gamma)\,,\qquad 0<s<1\,,\tag{6.2}$$

Therefore $\operatorname{dom}(\Theta_{N,\Sigma})\subseteq H^s(\Gamma)$, $\frac{1}{2}\le s<1$, and so, by Remark 4.7 and (3.13),

$$\operatorname{dom}(A_{N,\Sigma})\subseteq H^{\frac{3}{2}-}(\mathbb{R}^3\backslash\overline{\Sigma})\,.$$

6.3. *Robin boundary conditions.*

Given $\Sigma\subset\Gamma$ relatively open of class $C^{0,1}$, here we denote by $\Pi^{\oplus}_{\Sigma}$ the orthogonal projector in the Hilbert space $H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma)$ such that $\operatorname{ran}(\Pi^{\oplus}_{\Sigma})=H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^{\perp}\oplus H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^{\perp}$. By (3.4), $\operatorname{ran}((\Pi^{\oplus}_{\Sigma})')=\Lambda^{-3}H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)\oplus\Lambda^{-1}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$, where $(\Pi^{\oplus}_{\Sigma})'=(\Lambda^3\oplus\Lambda)\Pi^{\oplus}_{\Sigma}(\Lambda^{-3}\oplus\Lambda^{-1})$ is the dual projection. In the following, we also use the identifications $H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^{\perp}\oplus H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^{\perp}\simeq H^{\frac{3}{2}}(\Sigma)\oplus H^{\frac{1}{2}}(\Sigma)$ and $(H^{\frac{3}{2}}(\Sigma)\oplus H^{\frac{1}{2}}(\Sigma))'\simeq H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)\oplus H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$. We denote the orthogonal projection from $H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma)$ onto $H^{\frac{3}{2}}(\Sigma)\oplus H^{\frac{1}{2}}(\Sigma)$ by $R^{\oplus}_{\Sigma}$; it can be identified with the restriction map $R^{\oplus}_{\Sigma}:H^{s_1}(\Gamma)\oplus H^{s_2}(\Gamma)\to H^{s_1}(\Sigma)\oplus H^{s_2}(\Sigma)$, $R^{\oplus}_{\Sigma}(\phi\oplus\varphi):=(\phi|\Sigma)\oplus(\varphi|\Sigma)$ (see the end of Subsection 3.1).

Theorem 6.13. *Let Ω be of class $C^{4,1}$, and let $b_{\pm}\in M^{\frac{3}{2}}(\Gamma)$ be real valued and $1/[b]\in L^{\infty}(\Gamma)$ be negative. Then*

$$\Theta_{R,\Sigma}:\operatorname{dom}(\Theta_{R,\Sigma})\subseteq H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)\oplus H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)\to H^{\frac{3}{2}}(\Sigma)\oplus H^{\frac{1}{2}}(\Sigma)\,,$$

$$\begin{aligned}\Theta_{R,\Sigma}(\phi,\varphi):=&R^{\oplus}_{\Sigma}\begin{bmatrix}1/[b]+\gamma_0\mathrm{SL} & \langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL}\\ \langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL} & b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL}\end{bmatrix}(\Pi^{\oplus}_{\Sigma})'(\phi,\varphi)\\ \equiv&(((1/[b]+\gamma_0\mathrm{SL})\phi+(\langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL})\varphi)|\Sigma)\oplus((((\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\phi+(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma)\,,\end{aligned}$$

$$\begin{aligned}\operatorname{dom}(\Theta_{R,\Sigma}):=&\{(\phi,\varphi)\in L^2_{\overline{\Sigma}}(\Gamma)\times H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma):((1/[b]+\gamma_0\mathrm{SL})\phi+(\langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL})\varphi)|\Sigma\in H^{\frac{3}{2}}(\Sigma)\,,\\ &((\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\phi+(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma\in H^{\frac{1}{2}}(\Sigma)\}\end{aligned}$$

is self-adjoint.

Proof. Let $\tilde{\Theta}_R$ be the self-adjoint operator in $H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma)$

$$\tilde{\Theta}_R := \Theta_R(\Lambda^3\oplus\Lambda) : H^{9/2}(\Gamma)\times H^{5/2}(\Gamma)\subseteq H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma)\to H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma)\,.$$

Since $\gamma_0\mathrm{SL}\,\Lambda^3$ is infinitesimally $(1/[b])\Lambda^3$-bounded (proceed as in the proof of Lemma 5.3) and $b_+b_-/[b]\Lambda$ is infinitesimally $(\hat{\gamma}_1\mathrm{SL}\,\Lambda)$-bounded (proceed as in the proof of Lemma 5.4), by $[b]<0$ and (3.30) respectively, both $(1/[b]+\gamma_0\mathrm{SL})\Lambda^3$ and $(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\Lambda$ are upper bounded. Let λ_* be a common strict upper bound; by the Frobenius-Schur factorization of the block operator matrix $\tilde{\Theta}_R$ (see e.g. [66, Theorem 2.2.18]), there exists $\lambda_R>\lambda_*$ such that $-\tilde{\Theta}_R+\lambda_R>0$ whenever

$$-(1/[b]+\gamma_0\mathrm{SL})\Lambda^3+\lambda_R > (\langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL})\Lambda(-(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\Lambda+\lambda_R)^{-1}(\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^3\,. \tag{6.3}$$

Since

$$\langle(-(1/[b]+\gamma_0\mathrm{SL})\Lambda^3+\lambda_R)\phi,\phi\rangle_{H^{\frac{3}{2}}(\Gamma)}\ge(\lambda_R-\lambda_*)\,\|\phi\|^2_{H^{\frac{3}{2}}(\Gamma)}$$

and

$$\begin{aligned}&\langle((\langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL})\Lambda(-(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\Lambda+\lambda_R)^{-1}(\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^3\phi,\phi\rangle_{H^{\frac{3}{2}}(\Gamma)}\\ =&\langle(-(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\Lambda+\lambda_R)^{-1}(\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^3\phi,(\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^3\phi\rangle_{H^{\frac{1}{2}}(\Gamma)}\\ \le&\|(-(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\Lambda+\lambda_*)^{-1}\|_{H^{-\frac{1}{2}}(\Gamma),H^{\frac{3}{2}}(\Gamma)}\|(\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^{\frac{1}{2}}\|^2_{L^2(\Gamma),H^{-\frac{1}{2}}(\Gamma)}\|\phi\|^2_{H^{\frac{3}{2}}(\Gamma)}\,,\end{aligned}$$

the operator inequality (6.3) holds true by taking λ_R sufficiently large. Let f_R be the densely defined, semibounded, closed sesquilinear form associated with $\tilde{\Theta}_R$, i.e.

$$f_R : (H^3(\Gamma)\times H^{\frac{3}{2}}(\Gamma))\times(H^3(\Gamma)\times H^{\frac{3}{2}}(\Gamma))\subset(H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma))\times(H^{\frac{3}{2}}(\Gamma)\oplus H^{\frac{1}{2}}(\Gamma))\to\mathbb{R}\,,$$

$$\begin{aligned}f_R((\phi_1,\varphi_1),(\phi_2,\varphi_2)) :=&\langle(1/[b]+\hat{\gamma}_0\mathrm{SL})\Lambda^3\phi_1+(\langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL})\Lambda\varphi_1,\Lambda^3\phi_2\rangle_{L^2(\Gamma)}\\ &+\langle(\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^3\phi_1+(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\Lambda\varphi_1,\Lambda\varphi_2\rangle_{-\frac{1}{2},\frac{1}{2}}\,.\end{aligned}$$

Since $H^3(\Gamma)\cap H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp=H^3(\Gamma)\cap\Lambda^{-3}H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)=\Lambda^{-3}L^2_{\overline{\Sigma}}(\Gamma)$ is dense in $\Lambda^{-3}H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)$ and $H^{\frac{3}{2}}(\Gamma)\cap H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp=H^{\frac{3}{2}}(\Gamma)\cap\Lambda^{-1}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)=\Lambda^{-1}H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$ is dense in $\Lambda^{-1}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$, we can use Lemma 7.1 to determine the semibounded self-adjoint operator $\tilde{\Theta}_{R,\Sigma}$ in $\mathrm{ran}(\Pi^\oplus_\Sigma)$ associated to the restriction of f_R to $\mathrm{ran}(\Pi^\oplus_\Sigma)$. Then

$$\Theta_{R,\Sigma} := U^\oplus_\Sigma\tilde{\Theta}_{R,\Sigma}(\Lambda^{-3}\oplus\Lambda^{-1})\,,\quad \mathrm{dom}(\Theta_{R,\Sigma}) := (\Lambda^3\oplus\Lambda)\mathrm{dom}(\tilde{\Theta}_{R,\Sigma})\,,$$

is self-adjoint, where the map $U^\oplus_\Sigma\Pi^\oplus_\Sigma(\phi\oplus\varphi)=(\phi|\Sigma)\oplus(\varphi|\Sigma)$ provides the unitary isomorphism $(H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp\oplus H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp)\simeq(H^{\frac{3}{2}}(\Sigma)\oplus H^{\frac{1}{2}}(\Sigma))$. To conclude the proof we need to determine the operator $\breve{\Theta}_R := (\tilde{\Theta}_R)\breve{}$ and the subspace $K_\Sigma := \mathfrak{k}_{\Pi^\oplus_\Sigma}$ (we refer to the Appendix for the notations). Let H_R be the Hilbert space given by $\mathrm{dom}(f_R)=H^3(\Gamma)\times H^{\frac{3}{2}}(\Gamma)$ endowed with the scalar product

$$\langle(\phi_1,\varphi_1),(\phi_2,\varphi_2)\rangle_R := (-f_R+\lambda_R)((\phi_1,\varphi_1),(\phi_2,\varphi_2))\,.$$

Let H'_R denote its dual space. Since

$$\begin{aligned}&f_R((\phi_1,\varphi_1),(\phi_2,\varphi_2))=\langle(1/[b]+\hat{\gamma}_0\mathrm{SL})\Lambda^3\phi_1+(\langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL})\Lambda\varphi_1,\Lambda^3\phi_2\rangle_{L^2(\Gamma)}\\ &+\langle\Lambda^{-\frac{1}{2}}((\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^3\phi_1+(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL}))\Lambda\varphi_1,\Lambda^{\frac{3}{2}}\varphi_2\rangle_{L^2(\Gamma)}\,,\end{aligned}$$

one has

$$H'_R=L^2(\Gamma)\times H^{-\frac{1}{2}}(\Gamma)\,,\quad \langle(\phi',\varphi'),(\phi,\varphi)\rangle_{H'_R,H_R}=\langle\phi',\Lambda^3\phi\rangle_{L^2(\Gamma)}+\langle\Lambda^{-\frac{1}{2}}\varphi',\Lambda^{\frac{3}{2}}\varphi\rangle_{L^2(\Gamma)}\,,$$

and

$$\breve{\Theta}_R : H^3(\Gamma)\times H^{\frac{3}{2}}(\Gamma)\to L^2(\Gamma)\times H^{-\frac{1}{2}}(\Gamma)\,,$$

$$\breve{\Theta}_R(\phi,\varphi)=((1/[b]+\hat{\gamma}_0\mathrm{SL})\Lambda^3\phi+(\langle b\rangle/[b]+\hat{\gamma}_0\mathrm{DL})\Lambda\varphi,(\langle b\rangle/[b]+\hat{\gamma}_1\mathrm{SL})\Lambda^3\phi+(b_+b_-/[b]+\hat{\gamma}_1\mathrm{DL})\Lambda\varphi).$$

Moreover

$$\begin{aligned}K_\Sigma =&\{(\phi',\varphi')\in L^2(\Gamma)\times H^{-\frac12}(\Gamma) : \forall(\phi,\varphi)\in\Lambda^{-3}L^2_{\overline{\Sigma}}(\Gamma)\times\Lambda^{-1}H^{\frac12}_{\overline{\Sigma}}(\Gamma)\,,\ \langle(\varphi',\phi'),(\phi,\varphi)\rangle_{H_R',H_R}=0\}\\ =&\{(\phi',\varphi')\in L^2(\Gamma)\times H^{-\frac12}(\Gamma) : \forall\,(\phi,\varphi)\in L^2_{\overline{\Sigma}}(\Gamma)\times H^{\frac12}_{\overline{\Sigma}}(\Gamma)\,,\ \langle\phi',\phi\rangle_{L^2(\Gamma)}+\langle\varphi',\varphi\rangle_{-\frac12,\frac12}=0\}\\ =&L^2_{\Sigma^c}(\Gamma)\times H^{-\frac12}_{\Sigma^c}(\Gamma)\,.\end{aligned}$$

Therefore, by Lemma 7.1, $\tilde\Theta_{R,\Sigma}$ is self-adjoint on the domain

$$\begin{aligned}\mathrm{dom}(\tilde\Theta_{R,\Sigma}) :=&\{(\phi,\varphi)\in\Lambda^{-3}L^2_{\overline{\Sigma}}(\Gamma)\times\Lambda^{-1}H^{\frac12}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde\phi\oplus\tilde\varphi\in\mathrm{ran}(\Pi^\oplus_\Sigma)\text{ s.t.}\\ &(1/[b]+\hat\gamma_0\mathrm{SL})\Lambda^3\phi+(\langle b\rangle/[b]+\hat\gamma_0\mathrm{DL})\Lambda\varphi-\tilde\phi\in L^2_{\Sigma^c}(\Gamma)\text{ and}\\ &(\langle b\rangle/[b]+\hat\gamma_1\mathrm{SL})\Lambda^3\phi+(b_+b_-/[b]+\hat\gamma_1\mathrm{DL})\Lambda\varphi-\tilde\varphi\in H^{-\frac12}_{\Sigma^c}(\Gamma)\}\end{aligned}$$

and $\tilde\Theta_{R,\Sigma}(\phi,\varphi):=(\tilde\phi,\tilde\varphi)$. Then

$$\begin{aligned}&U^\oplus_\Sigma\tilde\Theta_{R,\Sigma}(\Lambda^{-3}\phi,\Lambda^{-1}\varphi)=(\tilde\phi|\Sigma)\oplus(\tilde\varphi|\Sigma)\\ =&(((1/[b]+\hat\gamma_0\mathrm{SL})\phi+(\langle b\rangle/[b]+\hat\gamma_0\mathrm{DL})\varphi)|\Sigma)\oplus(((\langle b\rangle/[b]+\hat\gamma_1\mathrm{SL})\phi+(b_+b_-/[b]+\hat\gamma_1\mathrm{DL})\varphi)|\Sigma)\end{aligned}$$

and

$$\begin{aligned}&\mathrm{dom}(\Theta_{R,\Sigma})\\ =&\{(\phi,\varphi)\in L^2_{\overline{\Sigma}}(\Gamma)\times H^{\frac12}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde\phi\oplus\tilde\varphi\in\mathrm{ran}(\Pi^\oplus_\Sigma)\text{ s.t.}\\ &(1/[b]+\hat\gamma_0\mathrm{SL})\phi+(\langle b\rangle/[b]+\hat\gamma_0\mathrm{DL})\varphi-\tilde\phi\in L^2_{\Sigma^c}(\Gamma)\text{ and}\\ &(\langle b\rangle/[b]+\hat\gamma_1\mathrm{SL})\phi+(b_+b_-/[b]+\hat\gamma_1\mathrm{DL})\varphi-\tilde\varphi\in H^{-\frac12}_{\Sigma^c}(\Gamma)\}\\ \subseteq&\{(\phi,\varphi)\in L^2_{\overline{\Sigma}}(\Gamma)\times H^{\frac12}_{\overline{\Sigma}}(\Gamma) : ((1/[b]+\hat\gamma_0\mathrm{SL})\phi+(\langle b\rangle/[b]+\hat\gamma_0\mathrm{DL})\varphi)|\Sigma\in H^{\frac32}(\Sigma)\text{ and}\\ &(((\langle b\rangle/[b]+\hat\gamma_1\mathrm{SL})\phi+(b_+b_-/[b]+\hat\gamma_1\mathrm{DL})\varphi)|\Sigma\in H^{\frac12}(\Sigma)\}\\ \subseteq&\{(\phi,\varphi)\in L^2_{\overline{\Sigma}}(\Gamma)\times H^{\frac12}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde\phi\oplus\tilde\varphi\in\mathrm{ran}(\Pi^\oplus_\Sigma)\text{ s.t.}\\ &((1/[b]+\hat\gamma_0\mathrm{SL})\phi+(\langle b\rangle/[b]+\hat\gamma_0\mathrm{DL})\varphi)|\Sigma=\tilde\phi|\Sigma\text{ and}\\ &(((\langle b\rangle/[b]+\hat\gamma_1\mathrm{SL})\phi+(b_+b_-/[b]+\hat\gamma_1\mathrm{DL})\varphi)|\Sigma=\tilde\varphi|\Sigma\}\\ =&\{(\phi,\varphi)\in L^2_{\overline{\Sigma}}(\Gamma)\times H^{\frac12}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde\phi\oplus\tilde\varphi\in\mathrm{ran}(\Pi^\oplus_\Sigma)\text{ s.t.}\\ &(1/[b]+\hat\gamma_0\mathrm{SL})\phi+(\langle b\rangle/[b]+\hat\gamma_0\mathrm{DL})\varphi-\tilde\phi\in L^2_{\Sigma^c}(\Gamma)\text{ and}\\ &(\langle b\rangle/[b]+\hat\gamma_1\mathrm{SL})\phi+(b_+b_-/[b]+\hat\gamma_1\mathrm{DL})\varphi-\tilde\varphi\in H^{-\frac12}_{\Sigma^c}(\Gamma)\}=\mathrm{dom}(\Theta_{R,\Sigma})\,.\end{aligned}$$

□

Corollary 6.14. *The linear operator in* $L^2(\mathbb{R}^n)$ *defined by* $A_{R,\Sigma}:=(A^{\max}_-\oplus A^{\max}_+)|\mathrm{dom}(A_{R,\Sigma})$*, where*

$$\begin{aligned}\mathrm{dom}(A_{R,\Sigma})=&\{u\in H^1(\mathbb{R}^n\backslash\overline{\Sigma})\cap(\mathrm{dom}(A^{\max}_-)\oplus\mathrm{dom}(A^{\max}_+)) : (-[\hat\gamma_1]u)\oplus[\gamma_0]u\in\mathrm{dom}(\Theta_{R,\Sigma})\,,\ (\gamma_1^\pm u-b_\pm\gamma_0^\pm u)|\Sigma=0\}\\ \subseteq&\{u\in H^1(\mathbb{R}^n\backslash\overline{\Sigma})\cap(\mathrm{dom}(A^{\max}_-)\oplus\mathrm{dom}(A^{\max}_+)) : (\gamma_1^\pm u-b_\pm\gamma_0^\pm u)|\Sigma=0\,,\ ([\hat\gamma_1]u)|\overline{\Sigma}^c=0\}\,,\end{aligned}$$

is self-adjoint and its resolvent is given by

$$(-A_{R,\Sigma}+z)^{-1}u=(-A+z)^{-1}u-G_z(\Pi^\oplus_\Sigma)'\left(R^\oplus_\Sigma\begin{bmatrix}1/[b]+\gamma_0\mathrm{SL}_z & \langle b\rangle/[b]+\gamma_0\mathrm{DL}_z\\ \langle b\rangle/[b]+\gamma_1\mathrm{SL}_z & b_+b_-/[b]+\gamma_1\mathrm{DL}_z\end{bmatrix}(\Pi^\oplus_\Sigma)'\right)^{-1}R^\oplus_\Sigma\begin{bmatrix}\gamma_0(-A+z)^{-1}u\\ \gamma_1(-A+z)^{-1}u\end{bmatrix},$$

where $R^\oplus_\Sigma$ *is the restriction operator* $R^\oplus_\Sigma(\phi\oplus\varphi)=(\phi|\Sigma)\oplus(\varphi|\Sigma)$*,* $(\Pi^\oplus_\Sigma)'$ *acts there as the inclusion map* $(\Pi^\oplus_\Sigma)' : H^{-\frac32}_{\overline{\Sigma}}(\Gamma)\oplus H^{-\frac12}_{\overline{\Sigma}}(\Gamma)\to H^{-\frac32}(\Gamma)\oplus H^{-\frac12}(\Gamma)$ *and* G_z *is defined in* (4.2).

Proof. By Theorem 6.1 and Theorem 4.4, taking $\Pi = \Pi^{\oplus}_{\Sigma}$ and $\Theta = -(U^{\oplus}_{\Sigma})^{-1}\Theta_{R,\Sigma}$, one gets the self-adjoint extension $(A^{\max}_{-} \oplus A^{\max}_{+})|\mathrm{dom}(A_{R,\Sigma})$ with domain (contained in $H^1(\mathbb{R}^n\backslash\Gamma)$ by $\mathrm{dom}(\Theta_{R,\Sigma}) \subseteq L^2(\Gamma)\times H^{\frac{1}{2}}(\Gamma)$ and Remark 4.7),

$$\begin{aligned}
&\mathrm{dom}(A_{R,\Sigma})\\
=&\{u \in H^1(\mathbb{R}^n\backslash\Gamma) \cap (\mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+})) : (-[\hat{\gamma}_1]u) \oplus [\gamma_0]u \in \mathrm{dom}(\Theta_{R,\Sigma})\,,\\
&\Pi^{\oplus}_{\Sigma}(\gamma_0(u + \mathrm{SL}[\hat{\gamma}_1 u] - \mathrm{DL}[\gamma_0 u]) \oplus \gamma_1(u + \mathrm{SL}[\hat{\gamma}_1 u] - \mathrm{DL}[\gamma_0 u])) = -(U^{\oplus}_{\Sigma}\Theta_{R,\Sigma})^{-1}((-[\hat{\gamma}_1]u) \oplus [\gamma_0]u)\}\\
=&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+})) : (-[\hat{\gamma}_1]u) \oplus [\gamma_0]u \in \mathrm{dom}(\Theta_{R,\Sigma})\,,\\
&(\gamma_0 u)|\Sigma + (\gamma_0\mathrm{SL}[\hat{\gamma}_1 u])|\Sigma - (\gamma_0\mathrm{DL}[\gamma_0 u])|\Sigma = ((1/[b] + \hat{\gamma}_0\mathrm{SL})[\hat{\gamma}_1]u - (\langle b\rangle/[b] + \hat{\gamma}_0\mathrm{DL})[\gamma_0]u)|\Sigma\\
&(\hat{\gamma}_1 u)|\Sigma + (\hat{\gamma}_1\mathrm{SL}[\hat{\gamma}_1 u])|\Sigma - (\hat{\gamma}_1\mathrm{DL}[\gamma_0 u])|\Sigma = ((\langle b\rangle/[b] + \hat{\gamma}_1\mathrm{SL})[\hat{\gamma}_1]u - (b_+b_-/[b] + \hat{\gamma}_1\mathrm{DL})[\gamma_0]u)|\Sigma\}\\
=&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+})) : (-[\hat{\gamma}_1]u) \oplus [\gamma_0]u \in \mathrm{dom}(\Theta_{R,\Sigma})\,,\\
&([b]\gamma_0 u - [\hat{\gamma}_1]u + \langle b\rangle[\gamma_0]u)|\Sigma = 0\,,\ ([b]\gamma_1 u - \langle b\rangle[\gamma_1]u + b_+b_-[\gamma_0]u)|\Sigma = 0\}\\
=&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_{-}) \oplus \mathrm{dom}(A^{\max}_{+})) :\\
&(-[\hat{\gamma}_1]u) \oplus [\gamma_0]u \in \mathrm{dom}(\Theta_{R,\Sigma})\,,\ (\gamma^{\pm}_1 u - b_{\pm}\gamma^{\pm}_0 u)|\Sigma = 0\}\,.
\end{aligned}$$

The formula giving $(-A_{R,\Sigma} + z)^{-1}$ is consequence of (4.7), since

$$\begin{aligned}
&\left(-U^{\oplus}_{\Sigma}\Theta_{R,\Sigma} + \Pi^{\oplus}_{\Sigma}\begin{bmatrix} \gamma_0(\mathrm{SL} - \mathrm{SL}_z) & \gamma_0(\mathrm{DL} - \mathrm{DL}_z) \\ \gamma_1(\mathrm{SL} - \mathrm{SL}_z) & \gamma_1(\mathrm{DL} - \mathrm{DL}_z) \end{bmatrix}(\Pi^{\oplus}_{\Sigma})'\right)^{-1}\Pi^{\oplus}_{\Sigma}\\
=&\left(-\Theta_{R,\Sigma} + R^{\oplus}_{\Sigma}\begin{bmatrix} \gamma_0(\mathrm{SL} - \mathrm{SL}_z) & \gamma_0(\mathrm{DL} - \mathrm{DL}_z) \\ \gamma_1(\mathrm{SL} - \mathrm{SL}_z) & \gamma_1(\mathrm{DL} - \mathrm{DL}_z) \end{bmatrix}(\Pi^{\oplus}_{\Sigma})'\right)^{-1}R^{\oplus}_{\Sigma}
\end{aligned}$$

and

$$-\Theta_{R,\Sigma} + R^{\oplus}_{\Sigma}\begin{bmatrix} \gamma_0(\mathrm{SL} - \mathrm{SL}_z) & \gamma_0(\mathrm{DL} - \mathrm{DL}_z) \\ \gamma_1(\mathrm{SL} - \mathrm{SL}_z) & \gamma_1(\mathrm{DL} - \mathrm{DL}_z) \end{bmatrix}(\Pi^{\oplus}_{\Sigma})' = R^{\oplus}_{\Sigma}\begin{bmatrix} 1/[b] + \gamma_0\mathrm{SL}_z & \langle b\rangle/[b] + \gamma_0\mathrm{DL}_z \\ \langle b\rangle/[b] + \gamma_1\mathrm{SL}_z & b_+b_-/[b] + \gamma_1\mathrm{DL}_z \end{bmatrix}(\Pi^{\oplus}_{\Sigma})'\,.$$

□

Remark 6.15. Since $\mathrm{supp}([\gamma_0]u) \subseteq \overline{\Sigma}$ and $\mathrm{supp}([\gamma_1]u) \subseteq \overline{\Sigma}$, for any $u \in \mathrm{dom}(A_{R,\Sigma})$, one has

$$A_{R,\Sigma}u = Au - [\gamma_1]u\,\delta_{\overline{\Sigma}} - [\gamma_0]u\,\partial_{\underline{a}}\delta_{\overline{\Sigma}} = Au - \frac{4}{[b]}\,\big((\langle b\rangle\,\gamma_1 u - b_+b_-\gamma_0 u)\,\delta_{\overline{\Sigma}} + (\gamma_1 u - \langle b\rangle\,\gamma_0 u)\,\partial_{\underline{a}}\delta_{\overline{\Sigma}}\big)$$

and so $(A_{R,\Sigma}u)|\overline{\Sigma}^c = (Au)|\overline{\Sigma}^c$. This also shows that $A_{R,\Sigma}$ is a self-adjoint extension of the symmetric operator $A|C^{\infty}_{\mathrm{comp}}(\mathbb{R}^n\backslash\overline{\Sigma})$. Hence it depends only on Σ and $b_{\pm}|\Sigma$ and not on the whole Γ: one would obtain the same operator by considering any other bounded domain $\Omega_{\circ}$ with boundary $\Gamma_{\circ}$ such that $\Sigma \subset \Gamma_{\circ}$.

Remark 6.16. By the results contained in the proof of Theorem 6.13, for the domain of the sesquilinear form $f_{R,\Sigma}$ associated to the self-adjoint operator $\tilde{\Theta}_{R,\Sigma}$ one has the relation $\mathrm{dom}(f_{R,\Sigma}) \subseteq H^3(\Gamma) \oplus H^{\frac{3}{2}}(\Gamma)$. Therefore Theorem 4.12 applies and so, by Corollary 4.13, $\sigma_{ac}(A_{R,\Sigma}) = \sigma_{ac}(A)$ and the wave operators $W_{\pm}(A_{R,\Sigma}, A)$, $W_{\pm}(A, A_{R,\Sigma})$ exist and are complete.

Remark 6.17. Let $\phi \oplus \varphi \in \mathrm{dom}(\Theta_{R,\Sigma})$. Then, by

$$((1/[b] + \gamma_0\mathrm{SL})\phi + (\langle b\rangle/[b] + \hat{\gamma}_0\mathrm{DL})\varphi)|\Sigma \in H^{\frac{3}{2}}(\Sigma)\,,$$

there follows $\phi|\Sigma \in H^{\frac{1}{2}}(\Sigma) \subseteq H^{s}_{\overline{\Sigma}}(\Gamma)$, $s < \frac{1}{2}$. Hence, by

$$((\langle b\rangle/[b] + \hat{\gamma}_1\mathrm{SL})\phi + (b_+b_-/[b] + \hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma)\,,$$

one gets $(\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma \in H^s(\Sigma)$, $s < \frac{1}{2}$. Using (4.6) and the same arguments as in the proof of Lemma 4.8, one obtains

$$\mathrm{dom}(A_{R,\Sigma}) \subseteq H^{2-}(\mathbb{R}^n\backslash(\overline{\Sigma} \cup (\partial\Sigma)^{\epsilon})) \cap H^2(\mathbb{R}^n\backslash\Sigma^{\epsilon})\,.$$

Remark 6.18. Suppose $n = 3$ and both Γ and Σ are smooth. By Theorem 6.13, we have

$$((\langle b\rangle/[b] + \hat{\gamma}_1\mathrm{SL})\phi + (b_+b_-/[b] + \hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma)$$

with $\phi \oplus \varphi \in \mathrm{dom}(\Theta_{R,\Sigma}) \subseteq L^2(\Gamma) \oplus H^{\frac{1}{2}}(\Gamma)$. This yields $(\hat{\gamma}_1\mathrm{DL}\varphi)|\Sigma \in L^2(\Sigma) \subseteq H^s(\Sigma)$, $s < 0$ and so, according to (6.2), we get $\varphi \in H^{s}_{\overline{\Sigma}}(\Gamma)$, $\frac{1}{2} \leq s < 1$. Therefore $\mathrm{dom}(\Theta_{R,\Sigma}) \subseteq L^2(\Gamma) \oplus H^s(\Gamma)$, $\frac{1}{2} \leq s < 1$, and so, by Remark 4.7 and (3.13),

$$\mathrm{dom}(A_{R,\Sigma}) \subseteq H^{\frac{3}{2}-}(\mathbb{R}^3\backslash\overline{\Sigma})\,.$$

6.4. *δ-interactions.*

Given $\Sigma \subset \Gamma$ relatively open of class $C^{0,1}$, we denote by Π_Σ the orthogonal projector in the Hilbert space $H^{\frac{3}{2}}(\Gamma)$ such that $\mathrm{ran}(\Pi_\Sigma) = H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp \simeq H^{\frac{3}{2}}(\Sigma)$. Here Π'_Σ, R_Σ and U_Σ denote the same operators as in Subsection 6.1.

Theorem 6.19. *Let Ω be of class $C^{4,1}$, let $\alpha \in M^{\frac{3}{2}}(\Gamma)$ be real-valued and let $1/\alpha \in L^\infty(\Gamma)$ have constant sign on (each connected component of) Γ. Then*

$$\Theta_{\alpha,D,\Sigma} : \mathrm{dom}(\Theta_{\alpha,D,\Sigma}) \subseteq H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{\frac{3}{2}}(\Sigma)\,,$$

$$\Theta_{\alpha,D,\Sigma}\phi := R_\Sigma(1/\alpha + \gamma_0\mathrm{SL})\Pi'_\Sigma\phi \equiv ((1/\alpha + \gamma_0\mathrm{SL})\phi)|\Sigma\,,$$

$$\mathrm{dom}(\Theta_{\alpha,D,\Sigma}) := \{\phi \in H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : ((1/\alpha + \gamma_0\mathrm{SL})\phi)|\Sigma \in H^{\frac{3}{2}}(\Sigma)\}\,,$$

is self-adjoint.

Proof. Since $\gamma_0\mathrm{SL}$ is infinitesimally $1/\alpha$-bounded (see the proof of Lemma 5.3), the self-adjoint operator in $H^{\frac{3}{2}}(\Gamma)$

$$\tilde\Theta_{\alpha,D} := \Theta_{\alpha,D}\Lambda^3 : H^{9/2}(\Gamma) \subseteq H^{\frac{3}{2}}(\Gamma) \to H^{\frac{3}{2}}(\Gamma)\,,$$

is semibounded: it is strictly positive whenever $1/\alpha \geq 0$ and upper bounded whenever $1/\alpha \leq 0$. In the following we suppose $1/\alpha \geq 0$, the case $1/\alpha \leq 0$ being treated in a similar way. Let $f_{\alpha,D}$ be the densely defined, strictly positive, closed sesquilinear form associated with $\tilde\Theta_{\alpha,D}$, i.e.

$$f_{\alpha,D} : H^3(\Gamma) \times H^3(\Gamma) \subseteq H^{\frac{3}{2}}(\Gamma) \times H^{\frac{3}{2}}(\Gamma) \to \mathbb{R}\,,$$

$$f_{\alpha,D}(\phi_1\phi_2) := \langle(1/\alpha + \gamma_0\mathrm{SL})\Lambda^3\phi_1, \Lambda^3\phi_2\rangle_{L^2(\Gamma)}\,.$$

Since $H^3(\Gamma) \cap H^{\frac{3}{2}}_{\Sigma^c}(\Gamma)^\perp = H^3(\Gamma) \cap \Lambda^{-3}H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) = \Lambda^{-3}L^2_{\overline{\Sigma}}(\Gamma)$ is dense in $\Lambda^{-3}H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma)$, we can use Lemma 7.1 to determine the semibounded self-adjoint operator $\tilde\Theta_{\alpha,D,\Sigma}$ in $\mathrm{ran}(\Pi_\Sigma)$ associated to the restriction of $f_{\alpha,D}$ to $\mathrm{ran}(\Pi_\Sigma)$. Then

$$\Theta_{\alpha,D,\Sigma} := U_\Sigma\tilde\Theta_{\alpha,D,\Sigma}\Lambda^{-3}\,, \quad \mathrm{dom}(\Theta_{\alpha,D,\Sigma}) := \Lambda^3\mathrm{dom}(\tilde\Theta_{\alpha,D,\Sigma})\,,$$

is self-adjoint. To conclude the proof we need to determine the operator $\breve\Theta_{\alpha,D} := (\tilde\Theta_{\alpha,D})\breve{}$ and the subspace $K_\Sigma := \mathfrak{k}_{\Pi_\Sigma}$ (we refer to the Appendix for the notations). Let $H_{\alpha,D}$ be the Hilbert space given by $\mathrm{dom}(f_{\alpha,D}) = H^3(\Gamma)$ endowed with the scalar product

$$\langle\phi_1, \phi_2\rangle_{\alpha,D} := f_{\alpha,D}(\phi_1, \phi_2)\,.$$

Let $H'_{\alpha,D}$ denote its dual space. Since

$$f_{\alpha,D}(\phi_1, \phi_2) = \langle(1/\alpha + \gamma_0\mathrm{SL})\Lambda^3\phi_1, \Lambda^3\phi_2\rangle_{L^2(\Gamma)}\,,$$

one has

$$H'_{\alpha,D} = L^2(\Gamma)\,, \quad \langle\varphi, \phi\rangle_{H'_{\alpha,D},H_{\alpha,D}} = \langle\varphi, \Lambda^3\phi\rangle_{L^2(\Gamma)}\,,$$

and

$$\breve\Theta_{\alpha,D} : H^3(\Gamma) \to L^2(\Gamma)\,, \quad \breve\Theta_{\alpha,D} = (1/\alpha + \gamma_0\mathrm{SL})\Lambda^3\,.$$

Moreover

$$\begin{aligned} K_\Sigma =& \{\varphi \in L^2(\Gamma) : \forall\phi \in \Lambda^{-3}L^2_{\overline{\Sigma}}(\Gamma)\,,\ \langle\varphi, \Lambda^3\phi\rangle_{L^2(\Gamma)} = 0\} \\ =& \{\varphi \in L^2(\Gamma) : \forall\phi \in L^2_{\overline{\Sigma}}(\Gamma)\,,\ \langle\varphi, \phi\rangle_{L^2(\Gamma)} = 0\} \\ =& L^2_{\Sigma^c}(\Gamma)\,. \end{aligned}$$

Therefore, by Lemma 5.1, $\tilde\Theta_{\alpha,D,\Sigma}$ is self-adjoint on the domain

$$\mathrm{dom}(\tilde\Theta_{\alpha,D,\Sigma}) := \{\phi \in \Lambda^{-3}L^2_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde\phi \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } (1/\alpha + \gamma_0\mathrm{SL})\Lambda^3\phi - \tilde\phi \in L^2_{\Sigma^c}(\Gamma)\}$$

and $\tilde\Theta_{\alpha,D,\Sigma}\phi := \tilde\phi$. Then

$$U_\Sigma\tilde\Theta_{\alpha,D,\Sigma}\Lambda^{-3}\phi = \tilde\phi|\Sigma = ((1/\alpha + \gamma_0\mathrm{SL})\phi)|\Sigma$$

and

$$\begin{aligned}\mathrm{dom}(\Theta_{\alpha,D,\Sigma}) =&\{\phi \in L^2_{\overline{\Sigma}}(\Gamma) : \exists\, \tilde\phi \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } (1/\alpha + \gamma_0\mathrm{SL})\phi - \tilde\phi \in L^2_{\Sigma^c}(\Gamma)\}\\ \subseteq&\{\phi \in L^2_{\overline{\Sigma}}(\Gamma) : ((1/\alpha + \gamma_0\mathrm{SL})\phi)|\Sigma \in H^{\frac{3}{2}}(\Sigma)\}\\ \subseteq&\{\phi \in L^2_{\overline{\Sigma}}(\Gamma) : \exists\, \tilde\phi \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } ((1/\alpha + \gamma_0\mathrm{SL})\phi)|\Sigma = \tilde\phi|\Sigma\}\\ =&\{\phi \in L^2_{\overline{\Sigma}}(\Gamma) : \exists\, \tilde\phi \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } (1/\alpha + \gamma_0\mathrm{SL})\phi - \tilde\phi \in L^2_{\Sigma^c}(\Gamma)\} = \mathrm{dom}(\Theta_{\alpha,D,\Sigma})\,.\end{aligned}$$

□

Remark 6.20. Let $\phi \in \mathrm{dom}(\Theta_{\alpha,D,\Sigma}) \subseteq L^2_{\overline{\Sigma}}(\Gamma)$. Then $\gamma_0\mathrm{SL}\phi \in H^1(\Gamma)$ and so, by $((1/\alpha + \gamma_0\mathrm{SL})\phi)|\Sigma \in H^{\frac{3}{2}}(\Sigma)$, one gets $(\phi/\alpha)|\Sigma \in H^1(\Sigma)$. Thus, since $H^s(\Sigma) = H^s_{\overline{\Sigma}}(\Gamma)$ for any $s \in [0, \frac{1}{2})$, in conclusion one has $\mathrm{dom}(\Theta_{\alpha,D,\Sigma}) \subseteq H^s(\Gamma)$ for any $s \in [0, \frac{1}{2})$.

Corollary 6.21. *The linear operator in $L^2(\mathbb{R}^n)$ defined by $A_{\alpha,\delta,\Sigma} := (A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{\alpha,\delta,\Sigma})$ with domain*

$$\begin{aligned}\mathrm{dom}(A_{\alpha,\delta,\Sigma}) =&\{u \in H^{2-}(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_1]u \in \mathrm{dom}(\Theta_{\alpha,D,\Sigma})\,,\ (\alpha\gamma_0 u - [\gamma_1]u)|\Sigma = 0\}\\ \subseteq&\{u \in H^{2-}(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : (\alpha\gamma_0 u - [\gamma_1]u)|\Sigma = 0\}\,,\end{aligned}$$

is self-adjoint and its resolvent is given by

$$(-A_{\alpha,\delta,\Sigma} + z)^{-1}u = (-A + z)^{-1} - \mathrm{SL}_z\Pi'_\Sigma(R_\Sigma(1 + \alpha\gamma_0\mathrm{SL}_z)\Pi'_\Sigma)^{-1}R_\Sigma\alpha\gamma_0(-A + z)^{-1}\,,$$

where R_Σ is the restriction operator $R_\Sigma\phi = \phi|\Sigma$ and Π'_Σ acts there as the inclusion map $\Pi'_\Sigma : H^{-\frac{3}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{-\frac{3}{2}}(\Gamma)$.

Proof. By Theorem 6.19 and Theorem 4.4, taking $\Pi(\phi \oplus \varphi) = \Pi_\Sigma\phi \oplus 0$ and $\Theta(\phi \oplus \varphi) = (-U^{-1}_\Sigma\Theta_{\alpha,D,\Sigma}\phi) \oplus 0$, one gets the self-adjoint extension $(A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{\alpha,\delta,\Sigma})$ with domain (contained in $H^s(\mathbb{R}^n\backslash\Gamma)$, $s < 2$, by $\mathrm{dom}(\Theta_{\alpha,D,\Sigma}) \subseteq H^s(\Gamma)$, $s < \frac{1}{2}$, and by Remark 4.7)

$$\begin{aligned}&\mathrm{dom}(A_{\alpha,\delta,\Sigma})\\ =&\{u \in H^{2-}(\mathbb{R}^n\backslash\Gamma) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_1]u \in \mathrm{dom}(\Theta_{\alpha,D,\Sigma})\,,\ \Pi_\Sigma(\gamma_0(u + \mathrm{SL}[\gamma_1 u])) = U^{-1}_\Sigma\Theta_{\alpha,D,\Sigma}[\gamma_1 u]\}\\ =&\{u \in H^{2-}(\mathbb{R}^n\backslash\Gamma) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_1]u \in \mathrm{dom}(\Theta_{\alpha,D,\Sigma})\,,\ (\gamma_0 u)|\Sigma + (\mathrm{SL}[\gamma_1 u])|\Sigma = ((1/\alpha + \gamma_0\mathrm{SL})[\hat\gamma_1 u])|\Sigma\}\\ =&\{u \in H^{2-}(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_1]u \in \mathrm{dom}(\Theta_{\alpha,D,\Sigma})\,,\ (\alpha\gamma_0 u - [\gamma_1]u)|\Sigma = 0\}\\ \subseteq&\{u \in H^{s}(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : (\alpha\gamma_0 u - [\gamma_1]u)|\Sigma = 0\}\,,\end{aligned}$$

The formula giving $(-A_{\alpha,\delta,\Sigma} + z)^{-1}$ is consequence of (4.7), since

$$(-U_\Sigma\Theta_{\alpha,D,\Sigma} + \Pi_\Sigma\gamma_0(\mathrm{SL} - \mathrm{SL}_z)\Pi'_\Sigma)^{-1}\Pi_\Sigma = (-\Theta_{\alpha,D,\Sigma} + R_\Sigma\gamma_0(\mathrm{SL} - \mathrm{SL}_z)\Pi'_\Sigma)^{-1}R_\Sigma$$

and

$$\begin{aligned}-\Theta_{\alpha,D,\Sigma}\phi + R_\Sigma\gamma_0(\mathrm{SL} - \mathrm{SL}_z)\phi =& -((1/\alpha + \gamma_0\mathrm{SL})\phi)|\Sigma + (\gamma_0\mathrm{SL}\phi)|\Sigma - (\gamma_0\mathrm{SL}_z\phi)|\Sigma\\ =& -((1/\alpha + \gamma_0\mathrm{SL}_z)\phi)|\Sigma = -(1/(\alpha|\Sigma))(1 + \alpha\gamma_0\mathrm{SL}_z)\phi)|\Sigma\,.\end{aligned}$$

□

Remark 6.22. Since $\mathrm{supp}([\gamma_0]u) \subseteq \overline{\Sigma}$, for any $u \in \mathrm{dom}(A_{\alpha,\delta,\Sigma})$, one has

$$A_{\alpha,\delta,\Sigma}\, u = Au - \alpha\gamma_0 u\, \delta_{\overline{\Sigma}}$$

and so $(A_{\alpha,\delta,\Sigma}\, u)|\overline{\Sigma}^c = (Au)|\overline{\Sigma}^c$. This also shows that $A_{\alpha,\delta,\Sigma}$ is a self-adjoint extension of the symmetric operator $A|C^\infty_{\mathrm{comp}}(\mathbb{R}^n\backslash\overline{\Sigma})$. Hence it depends only on Σ and $\alpha|\Sigma$ and not on the whole Γ: one would obtain the same operator by considering any other bounded domain $\Omega_\circ$ with boundary $\Gamma_\circ$ such that $\Sigma \subset \Gamma_\circ$.

Remark 6.23. According to Remark 6.20, $\phi|\Sigma \in H^1(\Sigma)$ whenever $\phi \in \mathrm{dom}(\Theta_{\alpha,D,\Sigma})$; thus Lemma 4.8 applies and one gets

$$\mathrm{dom}(A_{\alpha,\delta,\Sigma}) \subseteq H^2(\mathbb{R}^n\backslash(\overline{\Sigma} \cup (\partial\Sigma)^\epsilon))\,.$$

Remark 6.24. By the results contained in the proof of Theorem 6.19, for the domain of the sesquilinear form $f_{\alpha,D,\Sigma}$ associated to the self-adjoint operator $\tilde\Theta_{\alpha,D,\Sigma}$ one has the relation $\mathrm{dom}(f_{\alpha,D,\Sigma}) \subseteq H^3(\Gamma)$. Therefore Theorem 4.12 applies and so, by Corollary 4.13, $\sigma_{ac}(A_{\alpha,\delta,\Sigma}) = \sigma_{ac}(A)$ and the wave operators $W_\pm(A_{\alpha,\delta,\Sigma}, A)$, $W_\pm(A, A_{\alpha,\delta,\Sigma})$ exist and are complete.

6.5. δ'-interactions.

Given $\Sigma \subset \Gamma$ relatively open of class $C^{0,1}$, we denote by Π_Σ the orthogonal projector in the Hilbert space $H^{\frac{1}{2}}(\Gamma)$ such that $\operatorname{ran}(\Pi_\Sigma) = H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp \simeq H^{\frac{1}{2}}(\Sigma)$. Here Π'_Σ, R_Σ and U_Σ denote the same operators as in Subsection 6.2.

Theorem 6.25. *Let Ω be of class $C^{2,1}$ and let $\beta \in M^{\frac{1}{2}}(\Gamma)$ be real valued and $1/\beta \in L^\infty(\Gamma)$. Then*

$$\Theta_{\beta,N,\Sigma} : \operatorname{dom}(\Theta_{\beta,N,\Sigma}) \subseteq H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{\frac{1}{2}}(\Sigma)\,,$$

$$\Theta_{\beta,N,\Sigma}\phi := R_\Sigma(-1/\beta + \hat{\gamma}_1\mathrm{DL})\Pi'_\Sigma\phi \equiv (-1/\beta + \hat{\gamma}_1\mathrm{DL})\phi)|\Sigma\,,$$

$$\operatorname{dom}(\Theta_{\beta,N,\Sigma}) := \{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : ((-1/\beta + \hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma)\}\,,$$

is self-adjoint.

Proof. By (3.30), since $1/\beta$ is infinitesimally $\hat{\gamma}_1$DL-bounded (see the proof of Lemma 5.4), the self-adjoint operator in $H^{\frac{1}{2}}(\Gamma)$ given by

$$\tilde{\Theta}_{\beta,N} := \Theta_{\beta,N}\Lambda : H^{5/2}(\Gamma) \subseteq H^{\frac{1}{2}}(\Gamma) \to H^{\frac{1}{2}}(\Gamma)\,,$$

is upper bounded. Let $f_{\beta,N}$ be the densely defined, semibounded, closed sesquilinear form associated with $\tilde{\Theta}_{\beta,N}$, i.e.

$$f_{\beta,N} : H^{\frac{3}{2}}(\Gamma) \times H^{\frac{3}{2}}(\Gamma) \subset H^{\frac{1}{2}}(\Gamma) \times H^{\frac{1}{2}}(\Gamma) \to \mathbb{R}\,,$$

$$f_{\beta,N}(\varphi_1, \varphi_2) := \langle(-1/\beta + \hat{\gamma}_1\mathrm{DL})\Lambda\varphi_1, \Lambda\varphi_2\rangle_{-\frac{1}{2},\frac{1}{2}}\,.$$

Since $H^{\frac{3}{2}}(\Gamma) \cap H^{\frac{1}{2}}_{\Sigma^c}(\Gamma)^\perp = H^{\frac{3}{2}}(\Gamma) \cap \Lambda^{-1}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) = \Lambda^{-1}H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$ is dense in $\Lambda^{-1}H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)$, we can use Lemma 7.1 to determine the semibounded self-adjoint operator $\tilde{\Theta}_{\beta,N,\Sigma}$ in $\operatorname{ran}(\Pi_\Sigma)$ associated to the restriction of $f_{\beta,N}$ to $\operatorname{ran}(\Pi_\Sigma)$. Then

$$\Theta_{\beta,N,\Sigma} := U_\Sigma\tilde{\Theta}_{\beta,N,\Sigma}\Lambda^{-1}\,, \quad \operatorname{dom}(\Theta_{\beta,N,\Sigma}) := \Lambda\operatorname{dom}(\tilde{\Theta}_{\beta,N,\Sigma})\,,$$

is self-adjoint. To conclude the proof we need to determine the operator $\breve{\Theta}_{\beta,N} := (\tilde{\Theta}_{\beta,N})\breve{}$ and the subspace $K_\Sigma := \mathfrak{k}_{\Pi_\Sigma}$ (we refer to the Appendix for the notations). Let $H_{\beta,N}$ be the Hilbert space given by $\operatorname{dom}(f_{\beta,N}) = H^{\frac{3}{2}}(\Gamma)$ endowed with the scalar product

$$\langle(\varphi_1, \varphi_2\rangle_{\beta,N} := (-f_{\beta,N} + \lambda_{\beta,N})(\varphi_1\varphi_2)\,,$$

where $\lambda_{\beta,N}$ is chosen in such a way to have $-f_{\beta,N} + \lambda_{\beta,N} > 0$. Let $H'_{\beta,N}$ denote its dual space. Since

$$f_{\beta,N}(\varphi_1, \varphi_2) = \langle\Lambda^{-\frac{1}{2}}(-1/\beta - \hat{\gamma}_1\mathrm{DL}))\Lambda\varphi_1, \Lambda^{\frac{3}{2}}\varphi_2\rangle_{L^2(\Gamma)}\,,$$

one has

$$H'_{\beta,N} = H^{-\frac{1}{2}}(\Gamma)\,, \quad \langle(\phi, \varphi)\rangle_{H'_{\beta,N},H_{\beta,N}} = \langle\Lambda^{-\frac{1}{2}}\phi, \Lambda^{\frac{3}{2}}\varphi\rangle_{L^2(\Gamma)}\,,$$

and

$$\breve{\Theta}_{\beta,N} : H^{\frac{3}{2}}(\Gamma) \to H^{-\frac{1}{2}}(\Gamma)\,, \quad \breve{\Theta}_{\beta,N} = (-1/\beta + \hat{\gamma}_1\mathrm{DL})\Lambda\,.$$

Moreover

$$\begin{aligned} K_\Sigma =&\{\phi \in H^{-\frac{1}{2}}(\Gamma) : \forall\varphi \in \Lambda^{-1}H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)\,,\ \langle\Lambda^{-\frac{1}{2}}\phi, \Lambda^{\frac{3}{2}}\varphi\rangle_{L^2(\Gamma)} = 0\} \\ =&\{\phi \in H^{-\frac{1}{2}}(\Gamma) : \forall\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma)\,,\ \langle\phi, \varphi\rangle_{-\frac{1}{2},\frac{1}{2}} = 0\} \\ =&H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\,. \end{aligned}$$

Therefore, by Lemma 5.1, $\tilde{\Theta}_{\beta,N,\Sigma}$ is self-adjoint on the domain

$$\operatorname{dom}(\tilde{\Theta}_{\beta,N,\Sigma}) := \{\varphi \in \Lambda^{-1}H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde{\varphi} \in \operatorname{ran}(\Pi_\Sigma) \text{ s.t. } (-1/\beta + \hat{\gamma}_1\mathrm{DL})\Lambda\varphi - \tilde{\varphi} \in H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\}$$

and $\tilde{\Theta}_{\beta,N,\Sigma})\varphi := \tilde{\varphi}$. Then

$$U_\Sigma\tilde{\Theta}_{\beta,N,\Sigma}\Lambda^{-1}\varphi = \tilde{\varphi}|\Sigma = ((-1/\beta + \hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma$$

and

$$\begin{aligned}\mathrm{dom}(\Theta_{\beta,N,\Sigma}) =&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde{\varphi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } (-1/\beta + \hat{\gamma}_1\mathrm{DL})\varphi - \tilde{\varphi} \in H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\}\\ \subseteq&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : ((-1/\beta + \hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma)\}\\ \subseteq&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde{\varphi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } ((-1/\beta + \hat{\gamma}_1\mathrm{DL})\varphi)|\Sigma = \tilde{\varphi}|\Sigma\}\\ =&\{\varphi \in H^{\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) : \exists\,\tilde{\varphi} \in \mathrm{ran}(\Pi_\Sigma) \text{ s.t. } (-1/\beta + \hat{\gamma}_1\mathrm{DL})\varphi - \tilde{\varphi} \in H^{-\frac{1}{2}}_{\Sigma^c}(\Gamma)\} = \mathrm{dom}(\Theta_{\beta,N,\Sigma})\,.\end{aligned}$$

□

Corollary 6.26. *The linear operator in* $L^2(\mathbb{R}^n)$ *defined by* $A_{\beta,\delta',\Sigma} := (A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{\beta,\delta',\Sigma})$ *with domain*

$$\begin{aligned}\mathrm{dom}(A_{\beta,\delta',\Sigma}) =&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{\beta,N,\Sigma})\,,\ (\beta\hat{\gamma}_1 u - [\gamma_0]u)|\Sigma = 0\}\\ \subseteq&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : (\beta\gamma_0 u - [\hat{\gamma}_1]u)|\Sigma = 0\}\,,\end{aligned}$$

is self-adjoint and its resolvent is given by

$$(-A_{\beta,\delta',\Sigma} + z)^{-1}u = (-A + z)^{-1} + \mathrm{DL}_z\Pi'_\Sigma(R_\Sigma(1 - \beta\hat{\gamma}_1\mathrm{DL}_z)\Pi'_\Sigma)^{-1}R_\Sigma\beta\gamma_1(-A + z)^{-1}\,,$$

where R_Σ *is the restriction operator* $R_\Sigma\phi = \phi|\Sigma$ *and* Π'_Σ *acts there as the inclusion map* $\Pi'_\Sigma : H^{-\frac{1}{2}}_{\overline{\Sigma}}(\Gamma) \to H^{-\frac{1}{2}}(\Gamma)$.

Proof. By Theorem 6.19 and Theorem 4.4, taking $\Pi(\phi \oplus \varphi) = 0 \oplus \Pi_\Sigma$ and $\Theta(\phi \oplus \varphi) = 0 \oplus (-U_\Sigma^{-1}\Theta_{\beta,N,\Sigma}\phi)$, one gets the self-adjoint extension $(A^{\max}_- \oplus A^{\max}_+)|\mathrm{dom}(A_{\beta,\delta',\Sigma})$ with domain (contained in $H^1(\mathbb{R}^n\backslash\Gamma)$, by $\mathrm{dom}(\Theta_{\beta,N,\Sigma}) \subseteq H^{\frac{1}{2}}(\Gamma)$ and by Remark 4.7)

$$\begin{aligned}&\mathrm{dom}(A_{\beta,\delta',\Sigma})\\ =&\{u \in H^1(\mathbb{R}^n\backslash\Gamma) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{\beta,N,\Sigma})\,,\ \Pi_\Sigma(\gamma_1(u - \mathrm{DL}[\gamma_0 u])) = -U_\Sigma^{-1}\Theta_{\beta,N,\Sigma}[\gamma_0 u]\}\\ =&\{u \in H^1(\mathbb{R}^n\backslash\Gamma) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{\beta,N,\Sigma})\,,\ (\gamma_1 u)|\Sigma - (\mathrm{DL}[\gamma_0 u])|\Sigma = -((-1/\beta + \hat{\gamma}_1\mathrm{DL})[\gamma_0 u])|\Sigma\}\\ =&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : [\gamma_0]u \in \mathrm{dom}(\Theta_{\beta,N,\Sigma})\,,\ (\beta\hat{\gamma}_1 u - [\gamma_0]u)|\Sigma = 0\}\\ \subseteq&\{u \in H^1(\mathbb{R}^n\backslash\overline{\Sigma}) \cap (\mathrm{dom}(A^{\max}_-) \oplus \mathrm{dom}(A^{\max}_+)) : (\beta\hat{\gamma}_1 u - [\gamma_0]u)|\Sigma = 0\}\,,\end{aligned}$$

The formula giving $(-A_{\beta,\delta',\Sigma} + z)^{-1}$ is consequence of (4.7), since

$$(-U_\Sigma\Theta_{\beta,N,\Sigma} + \Pi_\Sigma\gamma_1(\mathrm{DL} - \mathrm{DL}_z)\Pi'_\Sigma)^{-1}\Pi_\Sigma = (-\Theta_{\beta,N,\Sigma} + R_\Sigma\gamma_1(\mathrm{DL} - \mathrm{DL}_z)\Pi'_\Sigma)^{-1}R_\Sigma$$

and

$$\begin{aligned}-\Theta_{\beta,N,\Sigma}\varphi + R_\Sigma\gamma_1(\mathrm{DL} - \mathrm{DL}_z)\varphi =&((1/\beta - \hat{\gamma}_1\mathrm{DL})\phi)|\Sigma + (\hat{\gamma}_1\mathrm{DL}\phi)|\Sigma - (\hat{\gamma}_1\mathrm{DL}_z\varphi)|\Sigma\\ =&((1/\beta - \hat{\gamma}_1\mathrm{DL}_z)\varphi)|\Sigma = (1/(\beta|\Sigma))(1 - \beta\hat{\gamma}_1\mathrm{DL}_z)\phi)|\Sigma\,.\end{aligned}$$

□

Remark 6.27. Since $\mathrm{supp}([\gamma_1]u) \subseteq \overline{\Sigma}$, for any $u \in \mathrm{dom}(A_{\beta,\delta',\Sigma})$, one has

$$A_{\beta,\delta',\Sigma}u = Au - \beta\gamma_1 u\,\partial_{\underline{a}}\delta_{\overline{\Sigma}}$$

and so $(A_{\beta,\delta',\Sigma}u)|\overline{\Sigma}^c = (Au)|\overline{\Sigma}^c$. This also shows that $A_{\beta,\delta',\Sigma}$ is a self-adjoint extension of the symmetric operator $A|C^\infty_{\mathrm{comp}}(\mathbb{R}^n\backslash\overline{\Sigma})$. Hence it depends only on Σ and $\beta|\Sigma$ and not on the whole Γ: one would obtain the same operator by considering any other bounded domain $\Omega_\circ$ with boundary $\Gamma_\circ$ such that $\Sigma \subset \Gamma_\circ$.

Remark 6.28. According to the definition of $\mathrm{dom}(\Theta_{\beta,N,\Sigma})$, one has $\mathrm{dom}(\Theta_{\beta,N,\Sigma}) = \mathrm{dom}(\Theta_{N,\Sigma})$; thus Lemma 4.8 applies and one gets

$$\mathrm{dom}(A_{\beta,\delta',\Sigma}) \subseteq H^2(\mathbb{R}^n\backslash(\overline{\Sigma} \cup (\partial\Sigma)^\epsilon))\,.$$

Remark 6.29. By the results contained in the proof of Theorem 6.25, for the domain of the sesquilinear form $f_{\beta,N,\Sigma}$ associated to the self-adjoint operator $\tilde{\Theta}_{\beta,N,\Sigma}$ one has the relation $\mathrm{dom}(f_{\alpha,N,\Sigma}) \subseteq H^{\frac{3}{2}}(\Gamma)$. Therefore Theorem 4.12 applies and so, by Corollary 4.13, $\sigma_{ac}(A_{\beta,\delta',\Sigma}) = \sigma_{ac}(A)$ and the wave operators $W_\pm(A_{\beta,\delta',\Sigma}, A)$, $W_\pm(A, A_{\beta,\delta',\Sigma})$ exist and are complete.

Remark 6.30. Let $\varphi \in \mathrm{dom}(\Theta_{\beta,N,\Sigma}) \subseteq H^{\frac{1}{2}}(\Gamma)$. Then $(\hat{\gamma}_1 \mathrm{DL}\varphi)|\Sigma \in H^{\frac{1}{2}}(\Sigma) \subseteq H^s(\Sigma)$, $s < 0$. Then, in case $n = 3$ and both Γ and Σ are smooth, by (6.2), $\varphi \in H^s(\Gamma)$, $s < 1$. Therefore $\mathrm{dom}(\Theta_{\beta,N,\Sigma}) \subseteq H^s(\Gamma)$, $\frac{1}{2} \le s < 1$, and so, by Remark 4.7 and (3.13),

$$\mathrm{dom}(A_{\beta,\delta',\Sigma}) \subseteq H^{\frac{3}{2}-}(\mathbb{R}^3 \backslash \overline{\Sigma}) \,.$$

7. Appendix. Some remarks on compressions of self-adjoint operators.

Let $\Theta : \mathrm{dom}(\Theta) \subseteq \mathfrak{h} \to \mathfrak{h}$ be a semibounded self-adjoint operator on the Hilbert space $\mathfrak{h}$ and let $f : \mathrm{dom}(f) \times \mathrm{dom}(f) \subseteq \mathfrak{h} \times \mathfrak{h} \to \mathbb{R}$ be the corresponding semibounded sesquilinear form. Without loss of generality, eventually by considering $(-\Theta)$ and/or adding a constant, we can suppose that Θ (and hence f) is strictly positive. Let $\mathfrak{h}_\Theta$ be the Hilbert space given by $\mathrm{dom}(f)$ endowed with the scalar product $\langle \varphi_1, \varphi_2 \rangle_{\mathfrak{h}_\Theta} := f(\varphi_1, \varphi_2)$. Let $\mathfrak{h}'_\Theta$ be its dual space and let $\iota : \mathfrak{h}_\Theta \to \mathfrak{h}'_\Theta$ be the injection defined by $\langle \iota\phi, \varphi \rangle_{\mathfrak{h}'_\Theta \mathfrak{h}_\Theta} = \langle \phi, \varphi \rangle_{\mathfrak{h}}$, where $\langle \cdot, \cdot \rangle_{\mathfrak{h}'_\Theta \mathfrak{h}_\Theta}$ denotes the $\mathfrak{h}'_\Theta$-$\mathfrak{h}_\Theta$ duality and $\langle \cdot, \cdot \rangle_{\mathfrak{h}}$ denotes the scalar product in $\mathfrak{h}$; by the identification $\iota\phi \equiv \phi$, we may regard $\mathfrak{h}_\Theta \subseteq \mathfrak{h} \subseteq \mathfrak{h}'_\Theta$. Let $\breve{\Theta} : \mathfrak{h}_\Theta \to \mathfrak{h}'_\Theta$ be the bounded operator defined by

$$\langle \breve{\Theta}\varphi_1, \varphi_2 \rangle_{\mathfrak{h}'_\Theta \mathfrak{h}_\Theta} = \langle \varphi_1, \varphi_2 \rangle_{\mathfrak{h}_\Theta} \,, \qquad \varphi_1, \varphi_2 \in \mathfrak{h}_\Theta \,.$$

Obviously $\breve{\Theta}|\mathrm{dom}(\Theta) = \Theta$; moreover, by [49, Theorem 2.1],

$$\mathrm{dom}(\Theta) = \mathfrak{D}_\Theta := \{\varphi \in \mathfrak{h}_\Theta : \breve{\Theta}\varphi \in \mathfrak{h}\} \,.$$

In conclusion one gets a well know characterization of Θ (see [34, Remark at page 13], [64, Proof of Theorem VIII.15]):

$$\Theta = \breve{\Theta}|\mathfrak{D}_\Theta \,. \tag{7.1}$$

Let now $\Pi : \mathfrak{h} \to \mathfrak{h}$ be an orthogonal projector such that $\mathrm{dom}(f) \cap \mathrm{ran}(\Pi)$ is dense in $\mathrm{ran}(\Pi)$. Then the sesquilinear form

$$f_\Pi : \mathrm{dom}(f_\Pi) \times \mathrm{dom}(f_\Pi) \subseteq \mathrm{ran}(\Pi) \times \mathrm{ran}(\Pi) \to \mathbb{R} \,,$$

$$f_\Pi(\varphi_1, \varphi_2) := f(\Pi\varphi_1, \Pi\varphi_2) = f(\varphi_1, \varphi_2) \,, \quad \mathrm{dom}(f_\Pi) := \mathrm{dom}(f) \cap \mathrm{ran}(\Pi) \,,$$

is densely defined, closed and strictly positive. Hence there exists a unique strictly positive self-adjoint operator Θ_Π in $\mathrm{ran}(\Pi)$ corresponding to f_Π. By the above reasonings applied to Θ_Π, we know that $\Theta_\Pi = \breve{\Theta}_\Pi|\mathfrak{D}_{\Theta_\Pi}$. For any $\varphi_1 \in \mathrm{dom}(\Theta) \cap \mathrm{ran}(\Pi)$ and for any $\varphi_2 \in \mathrm{dom}(f) \cap \mathrm{ran}(\Pi)$ one has

$$\langle \breve{\Theta}_\Pi\varphi_1, \varphi_2 \rangle_{\mathfrak{h}'_{\Theta_\Pi} \mathfrak{h}_{\Theta_\Pi}} = \langle \varphi_1, \varphi_2 \rangle_{\mathfrak{h}_{\Theta_\Pi}} = f(\Pi\varphi_1, \Pi\varphi_1) = \langle \Theta\Pi\varphi_1, \Pi\varphi_2 \rangle_{\mathfrak{h}} = \langle \Pi\Theta\Pi\varphi_1, \varphi_2 \rangle_{\mathfrak{h}} \,.$$

Thus $\mathrm{dom}(\Theta) \cap \mathrm{ran}(\Pi) \subseteq \mathfrak{D}_{\Theta_\Pi}$ and $\mathrm{C}_\Pi(\Theta) \subseteq \Theta_\Pi$, where $\mathrm{C}_\Pi(\Theta)$ is the compression of Θ to $\mathrm{ran}(\Pi)$ defined by

$$\mathrm{C}_\Pi(\Theta) : \mathrm{dom}(\mathrm{C}_\Pi(\Theta)) \subseteq \mathrm{ran}(\Pi) \to \mathrm{ran}(\Pi) \,,$$

$$\mathrm{dom}(\mathrm{C}_\Pi(\Theta)) := \mathrm{dom}(\Theta) \cap \mathrm{ran}(\Pi) \,, \quad \mathrm{C}_\Pi(\Theta)\phi := \Pi\Theta\Pi\phi = \Pi\Theta\phi \,.$$

Notice that $\mathrm{C}_\Pi(\Theta)$ is symmetric but it can be not self-adjoint; it is self-adjoint if and only if $\mathfrak{D}_{\Theta_\Pi} \subseteq \mathrm{dom}(\Theta)$.

We now give a more explicit definition of $\breve{\Theta}_\Pi$. Let $\mathfrak{h}_{\Theta_\Pi}$ be the Hilbert space $\mathrm{dom}(f_\Pi)$ endowed with the scalar product

$$\langle \varphi_1, \varphi_2 \rangle_{\mathfrak{h}_{\Theta_\Pi}} := f_\Pi(\varphi_1, \varphi_2) = f(\varphi_1, \varphi_2) = \langle \varphi_1, \varphi_2 \rangle_{\mathfrak{h}_\Theta}$$

and let $\mathfrak{h}'_{\Theta_\Pi}$ denote its dual. Since $\mathfrak{h}_{\Theta_\Pi} = \mathfrak{h}_\Theta \cap \mathrm{ran}(\Pi)$, by [6, Proposition 3.5.1] one has

$$\mathfrak{h}'_{\Theta_\Pi} = \mathfrak{h}'_\Theta / \mathfrak{k}_\Pi \,,$$

where

$$\mathfrak{k}_\Pi := \{\phi \in \mathfrak{h}'_\Theta : \forall\varphi \in \mathfrak{h}_{\Theta_\Pi} \,,\ \langle \phi, \varphi \rangle_{\mathfrak{h}'_\Theta \mathfrak{h}_\Theta} = 0 \,\} \,,$$

i.e.

$$\mathfrak{h}'_{\Theta_\Pi} = \{\, [\phi]\,,\ \phi \in \mathfrak{h}'_\Theta \}\,, \quad [\phi] := \{\psi \in \mathfrak{h}'_\Theta : \psi - \phi \in \mathfrak{k}_\Pi\}\,.$$

The $\mathfrak{h}'_{\Theta_\Pi}$-$\mathfrak{h}_{\Theta_\Pi}$ duality is then defined by $\langle [\phi], \varphi\rangle_{\mathfrak{h}'_{\Theta_\Pi}\mathfrak{h}_{\Theta_\Pi}} := \langle \phi, \varphi\rangle_{\mathfrak{h}'_\Theta \mathfrak{h}_\Theta}$. Let $\iota_\Pi : \mathfrak{h}_{\Theta_\Pi} \to \mathfrak{h}'_{\Theta_\Pi}$ be the injection defined by $\langle \iota_\Pi \phi, \varphi\rangle_{\mathfrak{h}'_{\Theta_\Pi}\mathfrak{h}_{\Theta_\Pi}} := \langle \phi, \varphi\rangle_{\mathfrak{h}_{\Theta_\Pi}}$. Since

$$\langle \phi, \varphi\rangle_{\mathfrak{h}_{\Theta_\Pi}} = \langle \phi, \varphi\rangle_{\mathfrak{h}_\Theta} = \langle \iota\phi, \varphi\rangle_{\mathfrak{h}'_\Theta \mathfrak{h}_\Theta} = \langle [\iota\phi], \varphi\rangle_{\mathfrak{h}'_{\Theta_\Pi}\mathfrak{h}_{\Theta_\Pi}}\,,$$

one gets $\iota_\Pi \varphi = [\iota\varphi]$. By the identification $\iota_\Pi\varphi \equiv \varphi$, we may regard $\mathfrak{h}_{\Theta_\Pi} \subseteq \operatorname{ran}(\Pi) \subseteq \mathfrak{h}'_{\Theta_\Pi}$. For any $\varphi_1, \varphi_2 \in \mathfrak{h}_{\Theta_\Pi}$, the bounded operator $\breve{\Theta}_\Pi : \mathfrak{h}_{\Theta_\Pi} \to \mathfrak{h}'_{\Theta_\Pi}$ satisfies the relations

$$\langle \breve{\Theta}_\Pi \varphi_1, \varphi_2\rangle_{\mathfrak{h}'_{\Theta_\Pi}, \mathfrak{h}_{\Theta_\Pi}} = f_\Pi(\varphi_1, \varphi_2) = f(\varphi_1, \varphi_2) = \langle \breve{\Theta}\varphi_1, \varphi_2\rangle_{\mathfrak{h}'_\Theta \mathfrak{h}_\Theta} = \langle [\breve{\Theta}\varphi_1], \varphi_2\rangle_{\mathfrak{h}'_{\Theta_\Pi}\mathfrak{h}_{\Theta_\Pi}}\,.$$

Thus $\breve{\Theta}_\Pi\varphi = [\breve{\Theta}\varphi]$ and

$$\mathfrak{d}_{\Theta_\Pi} = \{\varphi \in \mathfrak{h}_{\Theta_\Pi} : [\breve{\Theta}\varphi] \in \operatorname{ran}(\Pi)\} = \{\varphi \in \mathfrak{h}_{\Theta_\Pi} : \exists \tilde{\varphi} \in \operatorname{ran}(\Pi) \text{ s.t.}\, \breve{\Theta}\varphi - \tilde{\varphi} \in \mathfrak{k}_\Pi\}\,.$$

In conclusion we have the following

Lemma 7.1. *Let* $f : \operatorname{dom}(f) \times \operatorname{dom}(f) \subseteq \mathfrak{h} \times \mathfrak{h} \to \mathbb{R}$ *be the closed sesquilinear form corresponding to the semibounded self-adjoint operator* Θ*; let* $\Pi : \mathfrak{h} \to \mathfrak{h}$ *be an orthogonal projector such that* $\operatorname{dom}(f) \cap \operatorname{ran}(\Pi)$ *is dense in* $\operatorname{ran}(\Pi)$*. Then the self-adjoint operator* $\Theta_\Pi : \operatorname{dom}(\Theta_\Pi) \subseteq \operatorname{ran}(\Pi) \to \operatorname{ran}(\Pi)$ *associated to the closed semibounded sesquilinear form* f_Π *defined as the restriction of* f *to* $\operatorname{dom}(f) \cap \operatorname{ran}(\Pi)$*, is given by*

$$\operatorname{dom}(\Theta_\Pi) := \{\varphi \in \operatorname{dom}(f) \cap \operatorname{ran}(\Pi) : \exists \tilde{\varphi} \in \operatorname{ran}(\Pi) \text{ s.t.}\, \breve{\Theta}\varphi - \tilde{\varphi} \in \mathfrak{k}_\Pi\}\,, \qquad \Theta_\Pi\varphi := \tilde{\varphi}\,.$$

Moreover the compression $C_\Pi(\Theta)$ *is self-adjoint, equivalenty* $C_\Pi(\Theta) = \Theta_\Pi$*, if and only if* $\operatorname{dom}(\Theta_\Pi) \subseteq \operatorname{dom}(\Theta)$.

References

References

[1] M. S. Agranovich. *Sobolev spaces, their generalizations and elliptic problems in smooth and Lipschitz domains.* Springer, Berlin, 2015.

[2] S. Albeverio, J. Brasche, M. Röckner. Dirichlet forms and generalized Schrödinger operators. In: *Schrödinger operators*, 1-42, Lecture Notes in Phys., **345**, Springer, Berlin, 1989.

[3] S. Albeverio, J.E. Fenstad, R. H¯oegh-Krohn, W. Karwowski, T. Lindstr¯om. Perturbations of the Laplacian supported by null sets, with applications to polymer measures and quantum fields. *Phys. Lett. A*, **104**, 396-400, 1984.

[4] S. Albeverio, J.E. Fenstad, R. H¯oegh-Krohn, W. Karwowski, T. Lindstr¯om. Schrödinger operators with potentials supported by null sets, in *Ideas and Methods in Mathematical Analysis* (S. Albeverio, J. R. Fenstad, H. Holden, and T. Lindstrm, Eds.), Vol. II, Cambridge Univ. Press, New York/Cambridge, MA, 1991.

[5] J.-P. Antoine, F. Gesztesy, J. Shabani. Exactly solvable models of sphere interactions in quantum mechanics. *J. Phys. A* **20**, 3687-3712, 1987.

[6] J.P. Aubin. *Applied functional analysis.* 2nd edition, Wiley, New York, 1999.

[7] E. Balslev. The singular spectrum of elliptic differential operators in $L^p(\mathbb{R}^n)$. *Math. Scand.* **19**, 193-210, 1966.

[8] J. Behrndt, P. Exner, V. Lotoreichik. Schrödinger operators with δ-interactions supported on conical surfaces. J. Phys. A 47 (2014), no. 35, 355202, 16 pp.

[9] J. Behrndt, P. Exner, V. Lotoreichik. Schrödinger operators with δ- and δ'-interactions on Lipschitz surfaces and chromatic numbers of associated partitions. *Rev. Math. Phys.* **26**, 1450015, 43 pp., 2014.

[10] J. Behrndt, G. Grubb, M. Langer, V. Lotoreichik. Spectral asymptotics for resolvent differences of elliptic operators with δ and δ'-interactions on hypersurfaces. *J. Spectr. Theory* **5**, 697-729, 2015.

[11] J. Behrndt, M. Langer, V. Lotoreichik. Spectral Estimates for Resolvent Differences of Self-Adjoint Elliptic Operators. *Integr. Equ. Oper. Theory* **77**, 1-37, 2013

[12] J. Behrndt, M. Langer, V. Lotoreichik. Schrödinger operators with δ- and δ'-potentials supported on hypersurfaces. *Ann. Henri Poincaré* **14**, 385-423, 2013.

[13] J. Behrndt, T. Micheler. Elliptic differential operators on Lipschitz domains and abstract boundary value problems. *J. Funct. Anal.* **267**, 3657-3709, 2014.

[14] M.Sh. Birman, T.A. Suslina, R.G. Shterenberg. Absolute continuity of the two-dimensional Schrödinger operator with delta potential concentrated on a periodic system of curves. *Algebra i Analiz* **12**, 140-177, 2000, in Russian; translated in: *St. Petersburg Math. J.* **12**, 983-1012, 2001.

[15] J.F. Brasche, P. Exner, Yu. A. Kuperin. Schrödinger operators with singular interactions. *J. Math. Anal. Appl.* **184**, 112-139, 1994.

[16] J.F. Brasche, A. Teta. Spectral analysis and scattering theory for Schrödinger operators with an interaction supported by a regular curve. *Ideas and methods in quantum and statistical physics. Oslo,1988.* Cambridge Univ. Press, Cambridge, 197-211, 1992.

[17] J. Chazarain, A. Piriou. *Introduction to the theory of linear partial differential equations.* North-Holland, Amsterdam 1982.

[18] M. Costabel, E.P. Stephan. An Improved Boundary Galerkin Method for Three-Dimensional Crack Problems. *Integral Equations Operator Theory* **10**, 467-504, 1987.

[19] T. Coulhon, E. Russ, V. Tardivel-Nachef. Sobolev Algebras on Lie Groups and Riemannian Manifolds. *Am. J. Math.* **123**, 283-342, 2001.

[20] V. A. Derkach, M. M. Malamud. Generalized resolvents and the boundary value problems for Hermitian operators with gaps. *J. Funct. Anal.* **95**, 1-95, 1991.

[21] N. C. Dias, A. Posilicano, J. N. Prata. Self-adjoint, globally defined Hamiltonian operators for systems with boundaries. *Commun. Pure Appl. Anal.* **10**, 1687-1706, 2011

[22] P. Exner. Spectral properties of Schrödinger operators with a strongly attractive δ interaction supported by a surface. In: *Proc. NSF Summer Research Conference (Mt. Holyoke 2002)*, Contemp. Math., Amer. Math. Providence, R.I., 2003.

[23] P. Exner. Leaky quantum graphs: a review. In: *Analysis on Graphs and its Applications.* Selected papers based on the Isaac Newton Institute for Mathematical Sciences programme, Cambridge, UK, 2007. *Proc. Symp. Pure Math.* **77**, 523-564, 2008

[24] P. Exner, M. Praas. On geometric perturbations of critical Schrödinger operators with a surface interaction. *J. Math. Phys.* **50**, 112101, 2009.

[25] P. Exner, P., Ichinose, I.: Geometrically induced spectrum in curved leaky wires. *J. Phys. A* **34**, 1439-1450, 2001.

[26] P. Exner, S. Kondej. Curvature-induced bound states for a δ interaction supported by a curve in $\mathbb{R}^3$. Ann. Henri Poincaré **3**, 967-981, 2002.

[27] P. Exner, S. Kondej. Bound states due to a strong δ interaction supported by a curved surface. *J. Phys. A* **36**, 443-457, 2003.

[28] P. Exner, S. Kondej. Scattering by local deformations of a straight leaky wire. *J. Phys. A* **38**, 4865-4874, 2005.

[29] P. Exner, K. Nemcova. Leaky quantum graphs: approximations by point interaction Hamiltonians. J. Phys. A **36**, 10173-10193, 2003.

[30] P. Exner, K. Pankrashkin. Strong coupling asymptotics for a singular Schrödinger operator with an interaction supported by an open arc. *Comm. Partial Differential Equations*, **39**, 193-212, 2014.

[31] P. Exner, K. Yoshitomi. Band gap of the Schrödinger operator with a strong δ-interaction on a periodic curve. *Ann. Henri Poincaré* **2**, 1139-1158, 2001.

[32] P. Exner, K. Yoshitomi. Persistent currents for 2D Schrödinger operator with a strong δ-interaction on a loop. *J. Phys. A* **35**, 3479-3487, 2002.

[33] P. Exner, K. Yoshitomi. Eigenvalue asymptotics for the Schrödinger operator with a δ-interaction on a punctured surface. *Lett. Math. Phys.*, **65**, 19-26, 2003.

[34] W. G. Faris. *Self-Adjoint Operators*, Lecture Notes in Mathematics 433, Springer, Berlin, 1975

[35] R. Figari, A. Teta. Effective potential and fluctuations for a boundary value problem on a randomly perforated domain. *Lett. Math. Phys.* **26**, 295-305, 1992.

[36] R. Figari, A. Teta. A boundary value problem of mixed type on perforated domains. *Asymptotic Anal.* **6**, 271-284, 1993.

[37] F. Gesztesy, M. Mitrea. A description of all self-adjoint extensions of the Laplacian and Kreĭn-type resolvent formulas on non-smooth domains. *J. Anal. Math.* **113**, 53-172, 2011

[38] F. Gesztesy, D. Mitrea, I. Mitrea, M. Mitrea. On the nature of the Laplace-Beltrami operator on Lipschitz manifolds. *J. Math. Sci.* **172**, 279-346, 2011.

[39] P. Grisvard. *Elliptic problems in nonsmooth domains.* Pitman, Boston, 1985.

[40] G. Grubb. *Distributions and Operators.* Springer, New York, 2009.

[41] G. Grubb. Krein resolvent formulas for elliptic boundary problems in nonsmooth domains. *Rend. Semin. Mat. Univ. Politec. Torino* **66**, 271-297, 2008.

[42] J. Herczyński. On Schrödinger operators with distributional potentials. *J. Oper. Theory* **21**, 273-295, 1989.

[43] P. Hess, T. Kato. Perturbation of closed operators and their adjoints. *Comment. Math. Helv.* **45**, 524-529, 1970.

[44] L. Hörmander. The existence of wave operators in scattering theory. *Math. Z.* **146**, 69-91, 1976.

[45] L. Hörmander. *The Analysis of Linear Partial Differential Operators II.* Springer, Berlin 1983.

[46] G. C. Hsiao, W. L. Wendland. *Boundary Integral Equations.* Springer, Berlin 2008.

[47] G. Hu, A. Mantile, M. Sini. Direct and inverse acoustic scattering by a collection of extended and point-like scatterers. *Multiscale Model. Simul.*, **12**, 996-1027, 2014.

[48] T. Kato. Scattering theory with two Hilbert spaces. *J. Funct. Anal.* **1**, 342-369, 1967.

[49] T. Kato. *Perturbation Theory for Linear Operators.* Springer, Berlin 1976

[50] A. Kirsch, N. Grinberg. *The factorization method for inverse problems.* Oxford University Press, Oxford, 2008

[51] S. Kondej, I. Vegelic. Lower bounds on the lowest spectral gap of singular potential Hamiltonians. *Ann. Henri Poincaré* **8**, 109-134, 2007.

[52] M. V. Korovina. On the Schrödinger equation with potential concentrated on a surface. *Differ. Uravn.* **36**, 1115-1159, 2000, in Russian; translation in *Differ. Equ.* **36**, 1279-1283, 2000.

[53] I. Lobanov, V. Lotoreichik, I. Yu. Popov. A lower bound for the spectrum of the two-dimensional Schrödinger operator with a δ-potential on a curve. *Theor. Math. Phys.* **162**, 332-340, 2010.

[54] V. Lotoreichik. *Singular Values and Trace Formulae for Resolvent Power Differences of Self-Adjoint Elliptic Operators.* Monographic Series TU Graz, 2013

[55] M.M. Malamud. Spectral theory of elliptic operators in exterior domains. *Russ. J. Math. Phys.* **17**, 96-125, 2010.

[56] V.G. Maz'ya, T.O. Shaposhnikova. *Theory of Sobolev Multipliers.* Springer-Verlag, Berlin 2009

[57] W. McLean. *Strongly elliptic systems and boundary integral equations.* Cambridge University Press, Cambrdige, 2000.

[58] J-L. Lions E. Magenes. *Non-homogeneous Boundary Value Problems and Applications*. Springer, New York, 1972.

[59] B.S. Pavlov. Boundary conditions on thin manifolds and the semiboundedness of the threebody Schrödinger operator with point potential. *Math. Sb.* (N.S.), **136(178)**, 163-177, 1988, in Russian; translation in *Math. USSR-Sb.* **64**, 161-175, 1989.

[60] A. Posilicano. A Kreĭn-like formula for singular perturbations of self-adjoint operators and applications. *J. Funct. Anal.*, **183**, 109-147, 2001.

[61] A. Posilicano. Self-adjoint extensions by additive perturbations. *Ann. Sc. Norm. Super. Pisa Cl. Sci.*(5), **2**, 1-20, 2003.

[62] A. Posilicano. Boundary triples and Weyl functions for singular perturbations of self-adjoint operators. *Methods Funct. Anal. Topology*, **10**, 57-63, 2004.

[63] A. Posilicano. Self-adjoint extensions of restrictions, *Oper. Matrices*, **2**, 483–506, 2008.

[64] M. Reed and B. Simon, Methods of modern mathematical physics Vol. I: Functional Analysis. *Academic Press.* New York, 1972.

[65] K. Schmüdgen. *Unbounded self-adjoint operators on Hilbert space.* Springer, Dordrecht, 2012.

[66] C. Tretter: *Spectral theory of block operator matrices and applications.* Imperial College Press 2008.

[67] J. Wloka. *Partial Differential Equations.* Cambridge University Press 1987